

HERALD OF SCIENCE
OF S. SEIFULLIN KAZAKH AGROTECHNICAL
RESEARCH UNIVERSITY:
VETERINARY SCIENCES

№ 4 (008)

Astana 2024

COMPOSITION OF THE EDITORIAL BOARD

Editor in Chief:

Aitbay K. Bulashev - Doctor of Veterinary Sciences, Professor, S. Seifullin Kazakh Agrotechnical Research University, Republic of Kazakhstan, Astana

Members of the Editorial Board:

Altay E. Usenbaev - Candidate of Veterinary Sciences, Associate Professor, S. Seifullin Kazakh Agrotechnical Research University, Republic of Kazakhstan, Astana

Yersyn Y. Mukhanbetkaliyev - Candidate of Veterinary Sciences, Associate Professor, S. Seifullin Kazakh Agrotechnical Research University, Republic of Kazakhstan, Astana

Aida M. Abdibekova - Doctor of Veterinary Sciences, Professor, Kazakh Research Veterinary Institute, Republic of Kazakhstan, Almaty

Kanatbek N. Mukantayev - Doctor of Biological Sciences, National Center for Biotechnology, Republic of Kazakhstan, Astana

Orken S. Akibekov - Candidate of Veterinary Sciences, Associate Professor, S. Seifullin Kazakh Agrotechnical Research University, Republic of Kazakhstan, Astana

Akhan B. Abzhalelov - Doctor of Biological Sciences, Professor, L.N. Gumilev Eurasian National University, Republic of Kazakhstan, Astana

Elena V. Kukhar - Doctor of Biological Sciences, S. Seifullin Kazakh Agrotechnical Research University, Republic of Kazakhstan, Astana

Lyudmila A. Lider - Candidate of Veterinary Sciences, Associate Professor, S. Seifullin Kazakh Agrotechnical Research University, Republic of Kazakhstan, Astana

Balgabai S. Maikanov - Doctor of Biological Sciences, Professor, S. Seifullin Kazakh Agrotechnical Research University, Republic of Kazakhstan, Astana

Isatay T. Dzhakupov - Doctor of Veterinary Sciences, Professor, S. Seifullin Kazakh Agrotechnical Research University, Republic of Kazakhstan, Astana

Damegul T. Rakhimzhanova - Candidate of Veterinary Sciences, Associate Professor, S. Seifullin Kazakh Agrotechnical Research University, Republic of Kazakhstan, Astana

INTERNATIONAL EDITORIAL BOARD MEMBERS

Christian Matthias Bauer - Doctor of Veterinary Medicine, Professor, Justus Liebig University Giessen, Germany, Giessen

Ali Aydin - Professor, Istanbul University-Cerrahpaşa, Republic of Turkiye, Istanbul

Bu Zhigao - Professor, Harbin Veterinary Research Institute, People's Republic of China, Harbin

Krzysztof Anusz - PhD, professor, Warsaw University of Life Sciences, Poland, Warsaw

Ibrahim Che Omar - Professor, Universiti Malaysia Kelantan, Malaysia, Kota Bharu

Modestas Ruzauskas - Doctor of Medical Sciences, Professor, Affiliation Institute of Microbiology and Virology, Veterinary Academy, Lithuanian University of Health Sciences, Republic of Lithuania, Kaunas

Fedor I. Korenoy - Candidate of Geographical Sciences, M.V. Lomonosov Moscow State University, Russian Federation, Moscow

Tatiana I. Glotova - Doctor of Biological Sciences, Professor, Institute of Experimental Veterinary Medicine of Siberia and the Far East, Russian Federation, Novosibirsk

Maxim B. Rebezov - Doctor of Agricultural Sciences, Professor, V.M. Gorbатов Federal Scientific Center for Food Systems of the Russian Academy of Sciences, Russian Federation, Moscow

ISSN 2958-5430

eISSN 2958-5449

Index publications – 75830

Herald of Science of S.Seifullin Kazakh Agrotechnical Research University: Veterinary Sciences.
– Astana: S. Seifullin Kazakh Agrotechnical Research University, 2024. – № 4 (008). – P. 4-15.
- ISSN 2958-5430, ISSN 2958-5449










VETERINARY SCIENCES

doi.org/ 10.51452/kazatuvc.2024.4(008).1780

UDC 576.8

Research article

Study of the Immunologic Value of *Trichinella spiralis* recombinant Protein Tsp-LE

Orken S. Akibekov¹ , Fabio Tosini² , Alfiya S. Syzdykova³ , Aissarat M. Gajimuradova³ ,
Dariga D. Zhumalina³ , Fariza S. Zhagipar³ , Nurtai N. Gubaidullin³ ,
Nasipkhan A. Askarova³ , Saltanat S. Yesseneyeva⁴ 

¹Department of Microbiology and Biotechnology, Faculty of Veterinary and Livestock Technology,
S. Seifullin Kazakh Agrotechnical Research University, Astana, Kazakhstan

²European Union Reference Laboratory for Parasites, «Istituto Superiore di Sanità», Roma, Italy

³Research platform for agricultural biotechnology,

S. Seifullin Kazakh Agrotechnical Research University, Astana, Kazakhstan

⁴Department of Veterinary Sanitation, Faculty of Veterinary and Livestock Technology,
S. Seifullin Kazakh Agrotechnical Research University, Astana, Kazakhstan

Corresponding author: Orken S. Akibekov: orken.a.s@mail.ru

Co-authors: (1: FT) fabio.tosini@iss.it; (2: AS) halik.kz@mail.ru; (3: AG) aisarat3878@mail.ru;

(4: DZh) dariga.ast@gmail.com; (5: FZh) zhagipar.fariza@gmail.com;

(6: NG) nur-tai.kz@mail.ru; (7: NA) nasiko.askarova@bk.ru; (8: SYe) eseneeva.saltanat@mail.ru

Received: 01-10-2024 **Accepted:** 26-12-2024 **Published:** 30-12-2024

Abstract

Background and Aim. Trichinellosis is a serious zoonotic disease, with frequent outbreaks reported across various regions in Europe and Asia. However, current diagnostic tests based on excretory-secretory antigens (ES-Ag) and somatic antigens (S-Ag) lack adequate specificity and sensitivity. Moreover, the production of these antigens is both labor-intensive and costly.

Materials and Methods. The immunodiagnostic potential of a recombinant 11 kDa serine protease protein, previously characterized in earlier studies was assessed. This study detected infection in *Balb/c* mice as early as 14 days post-infection with *T. spiralis* larvae, yielding no false-positive results and an average infection level of 61±5.3 larvae per mouse.

Results. Comparative indirect ELISA analysis on mouse sera after the 3rd and 5th immunizations, alongside tests on S-Ag and S-Ag using sera from infected rabbits and pigs, demonstrated the high diagnostic accuracy of the recombinant protein. The protein revealed strong specificity for trichinellosis detection, showing no cross-reactivity with echinococcosis.

Conclusion. These findings support using rTsp-LE as a reliable component in developing ELISA test systems for trichinellosis detection.

Keywords: ELISA; Es-Ag; immunization; recombinant protein; S-Ag; *Trichinella spiralis*.

Introduction

Trichinellosis is a zoonotic disease caused by nematodes of the family *Trichinellidae*, affecting humans and a range of animals. The disease is registered in domestic animals such as pigs, dogs, and cats, as well as in numerous wild carnivores (including wolves, foxes, and bears), rodents (such as rats and mice), and humans [1]. Approximately 11 million people worldwide are estimated to be chronically infected with *Trichinella spiralis* [2].

In 2020, 181 cases of trichinellosis were reported across nine EU/EEA countries, with 117 cases confirmed and 64 classified as probable [3]. In China, from 2009 to 2020, cases of human trichinellosis have been primarily concentrated in southwestern regions, with eight outbreaks resulting in 479 cases and two fatalities [4]. Southeast Asia also shows variable seroprevalence rates of human trichinellosis, particularly in Laos and Vietnam, where rates range from 0% to 10.5% in certain villages. Additionally, in Cambodia, Laos, Malaysia, Thailand, and Vietnam, 13 outbreaks involving 1,604 cases and a mortality rate of less than 1% have been recorded over the past 21 years [5].

Given the epidemiological context, effective early diagnosis of trichinellosis in potentially high-risk domestic and wild animals is essential. However, routine testing of meat and animals for trichinellosis remains limited, primarily due to inflated costs, which can reach up to \$3 per animal [6].

Early clinical diagnosis of trichinellosis is challenging due to the lack of pronounced clinical symptoms. Moreover, diagnosing the disease in its chronic stages proves even more difficult [7].

The detection of anti-*Trichinella* IgG via enzyme-linked immunosorbent assay (ELISA) using muscle excretory-secretory antigens (Es-Ag) from *Trichinella spiralis* larvae is the most used serologic method for diagnosing trichinellosis. However, a major limitation of current serologic tests is the potential for false negatives during early infection stages [8, 9, 10]. Immunoenzyme methods utilizing recombinant antibodies promotes the diagnostic process, eliminating the need for animal slaughter, reducing reliance on antigen availability, standardizing components, and enhancing both the quality and specificity of ELISA-based diagnostics [11, 12].

Sufficient literature supports the high sensitivity and specificity of ELISA with recombinant antigens. For instance, Wang et al. (2015) studied a recombinant 31 kDa protein and found a sensitivity of 96.67% and 96.87%, compared to 100% and 98.44% for ELISA using ES antigens. ELISA with recombinant antigens was able to detect antibodies in mice with heavy (500 larvae), moderate (300 larvae), and light (100 larvae) infections at 8, 12, and 14 days post-infection (dpi), respectively, whereas detection using ES antigens occurred at 10, 8, and 10 dpi, respectively [13]. In mice infected with 200 muscle larvae (ML) of *T. spiralis*, the rTsTryp-ELISA demonstrated a natural muscle protease sensitivity of 98.1%, compared to 94.2% for the ES antigen-ELISA. Additionally, the specificity of rTsTryp-ELISA was significantly higher than that of the ES antigen-ELISA, at 98.7% versus 95.4%, respectively [12]. Somatic antigens with molecular masses of 43 kDa (TsCSAg-43), 79 kDa (TsCSAg-79), and 101 kDa (TsCSAg-101), isolated by Supcharoengoon et al. (2022) from the sera of trichinellosis-infected pigs, showed IgG-ELISA sensitivities of 100%, with specificities of 97.77%, 95.54%, and 90.63%, respectively [14].

Y. Liu et al. (2021) developed a novel competitive enzyme-linked immunoassay (rCLP-cELISA) based on recombinant cystatin-like protein and monoclonal antibodies. This method demonstrated high sensitivity and specificity with human serum field samples and showed robust applicability with porcine and murine serum [15]. The expression of *Trichinella* antigens encoding proteins of 53 kDa [16, 17], 49 kDa [18, 13], 35.5 kDa [19], and 21 kDa [20] in heterologous systems has become a standard approach for generating antigens used in diagnostic test systems.

Recombinant antigens maintain great promise for several reasons: they exhibit high specificity, minimizing the risk of false negatives and cross-reactions. Also, the expression of recombinant antigens enables standardized production with high product purity and stability, while also reducing the risk of infection for personnel handling parasites.

The aim of our study was to evaluate the immunodiagnostic potential of recombinant *Trichinella spiralis* serine protease protein at various stages of infection in experimentally infected laboratory mice.

Materials and Methods

Ethical approval. All animal procedures were conducted following exacting standards of biosafety and animal welfare. Protocols adhered to the International Guidelines for Biomedical Research Involving Animals [21].

The care and use of laboratory animals were approved by the Animal Ethics Committee of the Faculty of Veterinary Medicine and Animal Husbandry Technology at S. Seifullin Kazakh Agrotechnical Research University (KATRU), Astana, Kazakhstan (Protocol No. 1, dated February 23, 2023).

Experiments were conducted at the Research Platform of Agricultural Biotechnology, NCJSC: “S. Seifullin Kazakh Agrotechnical Research University.”

Infestation of animals *T. spiralis* larvae were maintained through serial passage in Soviet Chinchilla breed rabbits at the Joint Kazakh-Chinese Laboratory of Biological Safety.

Larvae were collected through artificial digestion following the standard protocol described previously [22]. For each species of trichinellosis, seven mice, aged three to four months, were selected for the experiment at 3, 5, 7-, 14-, 21-, and 30-days post-infection, with three mice forming the control group. Using the principle of analogs for *T. spiralis*, an experimental group of three mice was formed for each day of selection (18 in total), along with three control mice. The infection dose was 250 larvae per mouse, and the animals were orally infected using a disposable pipette containing *Trichinella larvae*.

Scheme of experiment. On the 3rd, 5th, 7th, 14th, 21st, and 30th days, three mice from each group were euthanized for pathological autopsy. Euthanasia was performed via sequential intramuscular injection of xylazine (Bioveta, Czech Republic) at a dose of 1.5 mg/kg, followed by intravenous injection of Anestofol (‘VIK’ LLC, Russia) at a dose of 7.5 mg/kg. Muscles were examined for the presence of parasites according to the guidelines of H.R. Gamble et al. (2019) [23]. The small intestine was dissected longitudinally, washed three times with ice-cold saline solution, then cut into 2-cm fragments and cultured in saline at 37 °C for 2.5 hours. Larvae that emerged from the small intestine into the physiological solution were collected using the Berman method [24].

Larvae isolation. *Trichinella* larvae were isolated from animal muscle tissue samples using compressor trichinelloscopy and digestion in artificial gastric juice (AGJ), following the methods outlined in Methodological Guidelines 4.2.2747-10 “Methods of Sanitary-Parasitological Examination of Meat and Meat Products” [25]. The detected and isolated helminthological material was preserved in a 70% ethanol solution.

Antiserum testing by indirect ELISA. ELISA was performed as previously described [13]. Briefly, 96-well ELISA plates (Corning, USA) were coated overnight at 4 °C with 100 µL of purified recombinant antigens (1 µg/mL) or ES antigens (1 µg/mL) in bicarbonate buffer (pH 9.6). After blocking with PBS containing 0.1% BSA for 1 hour at 37 °C, the following reagents were added sequentially and incubated for 1 hour at 37 °C: (1) mouse serum diluted 1:100 in PBST, and (2) HRP-conjugated anti-mouse IgG (Sigma, USA) diluted 1:5000. Reactions were detected by adding a one-component TMB substrate (tetramethylbenzidine, CJSC “HBO Immunotech”, Russia), and the reaction was stopped by adding 50 µL/well of 2M H₂SO₄. Optical density (OD) values at 450 nm were measured using a microplate reader (BioSan, Lithuania). All samples were analyzed in duplicate. A ratio of test sample OD to negative control OD (S/N ratio) < 2.1 was considered negative, while S/N ≥ 2.1 was considered positive.

Statistical analysis. Mean and standard deviation were calculated using Microsoft Excel 2010. For each value, Student's *t* was calculated, and a *p*-value < 0.05 was considered statistically significant. Serological studies were carried out using the method described by T.S. Saiduldin (1981) [26].

Results

During experiments infecting laboratory Balb/c mice with *T. spiralis* larvae at a dose of 250 larvae, the phase of larval migration from the intestine to the muscle was observed. The results indicate that larval migration begins as early as day 14, which is consistent with multiple studies. Figure 1 shows the numerical values of larvae detected through postmortem examination of the mice.





Figure 1 – Development of trichinella larvae in intestine and muscle

As shown in Figure 1, larvae were first detected in the intestines of infected mice at 7 dpi. However, by 14 dpi, active migration of *T. spiralis* larvae to the muscle was observed, with an average of 61 ± 5.3 larvae. A significant decrease in the number of larvae in the intestine was noted by 21 dpi (14 ± 2.08) ($p < 0.01$), accompanied by a 34.4% increase in muscle larvae ($p < 0.05$). By 30 dpi, the number of muscle larvae reached 110 ± 4.3 ($p < 0.01$), while the number of intestinal larvae decreased to 1.6 ± 0.88 ($p < 0.01$). Figure 2 shows muscle larvae isolated at 21 days post-infection.

On day 14, muscle larvae found in the muscles were both spirally twisted and hairpin-shaped, characteristics typical of the early muscle stage. This suggests ongoing larval migration from the intestine. Figure 2 shows muscle larvae isolated at 21 days post-infection.



T. spiralis larvae at 21 dpi found in mouse intestines

T. spiralis larvae at 21 dpi found in mouse muscles

Figure 2 – *Trichinella* larvae in the muscles of BALB/c mice (21 dpi)

Blood serum was collected from mice at each stage of infestation for antibody analysis using indirect ELISA. The results are shown in Table 1.

Table 1 – Antibody titer of mice infected with <i>T.spiralis</i> against ES-Ag measured by indirect ELISA Inventory /number of mice	Days after invasion of animals						Mean antibody titers
	3 days	5 days	7 days	14 days	21 days	30 days	
1	RA	RA	RA	1:800	1: 1600	1:3200	1:1600 (+18.9; -15.9)
2	RA	RA	1:100	1:800	1:3200	1:6400	1:3200 (+20.6; -17.1)
3	RA	RA	RA	1:400	1:1600	1:3200	1:980 (+12.5; -11.1)
Control group							
4	RA	RA	RA	RA	RA	RA	-
5	RA	RA	RA	RA	RA	RA	-
6	RA	RA	RA	RA	RA	RA	-

Note: Reaction absence

The tested method showed low efficiency in detecting trichinellosis during the intestinal phase of infection, despite the presence of larvae in the intestine. Antibodies were first detected at 14 days post-infection (dpi) with titers of 1:400 to 1:800 in some mice within the group. By day 30, titers had increased to 1:3200 and 1:6400, confirming a positive result. However, results obtained between days 7 and 14 could be interpreted as either positive or potentially false positive.

The recombinant serine protease protein Tsp-LE 11 kDa (PP099881.1), based on the cDNA sequence of the large exon of the *T. spiralis* serine protease gene (data not shown), was also evaluated for its diagnostic value. Studies were conducted on mouse, rabbit, and pig sera using Es-Ag, S-Ag, and the test antigen rTsp-LE.

An initial ELISA was performed using sera from mice immunized with the recombinant protein. The results are summarized in Table 2.

Table 2 – ELISA results using Es-Ag and rTsp-LE antigens in mice following two consecutive immunizations

Antigen	Average antibody titer	
	after the 3 rd immunization	after the 5 th immunization
ES-AG	1:300 (+41.4; -29.3)	1:570 (+23.1; -18.7)
Recombinant protein	1:570 (+23.1; -18.7)	1:2260 (+14.9; -13.0)
P	0.002	0.0001

During the first two immunizations, specific IgG was first detected in the infected mice serum after the third immunization with S-Ag, with OD 450 values of 0.324 and 0.234 at dilutions of 1:200 and 1:400, respectively. When using the recombinant antigen, a positive result was also observed after the third immunization at nearly the same titers of 1:400-1:800, with OD 450 values of 0.277 and 0.205, respectively. After the fifth immunization, detection occurred at titers of 1:400-1:800 and 1:1600-1:3200, with mean OD 450 values of 0.326 and 0.231 for ES-Ag and rTsp-LE, respectively. Despite the high specificity of ES-Ag, the ELISA values for detecting *trichinellosis* in mice using rTsp-LE are only slightly lower. However, it should be noted that the 11 kDa recombinant protein binds specifically

to the serine protease protein of *Trichinella* larvae, as evidenced by the absence of cross-reaction with *Echinococcus*, a commonly detected disease. Furthermore, this antigen was successfully used to diagnose animals infected with *T. nativa*, yielding results analogous to those obtained with *T. spiralis* (1:3200).

To study species specificity and the reaction of the obtained protein with antibodies from other animal species, indirect ELISA was performed using S-Ag, Es-Ag, and the recombinant protein. The results are presented in the table below.

Table 3 – Determination of antibody titer in sera of infected rabbits and pigs using Es-Ag, S-Ag and rTsp-LE

Antigens	Average antibody titer	
	Infested rabbits	Positive swine serum
S-Ag	1:20 480 (+14.1; -12.3)	1:6 400 (+32.0; -24.2)
ES-Ag	1:20 480 (+14.1; -12.3)	1:3 940 (+39.5; -28.2)
Recombinant protein	1:2 260 (+14.9; -13.0)	1:800 (+41.4; -29.3)
P	0.003	0.001

When sera from infected rabbits and pigs were tested using the three types of antigens, S-Ag showed a positive reaction with a mean titer of 1:20,480 in rabbits and 1:6,400 in pigs, with mean OD 450 values of 0.631 and 0.332, respectively. The results for ES-Ag were nearly identical to those for S-Ag, with titers of 1:20,480 in rabbits and 1:3,940 in pigs, and OD 450 values of 0.581 and 0.456, respectively. The recombinant Tsp-LE protein yielded positive results at mean titers of 1:2,600 in rabbits and 1:800 in pigs, with mean OD 450 values of 0.225 and 0.226, respectively.

Based on the results, the recombinant 11 kDa rTsp-LE protein demonstrates sufficient diagnostic value for detecting trichinellosis infection in different animal species. This protein can be utilized for the further development of a highly specific ELISA test.

Discussion and Conclusion

ES antigens derived from the muscle tissue of *T. spiralis* larvae are widely used as serodiagnostic agents for *trichinellosis* and are recommended by the International Commission on Trichinellosis (ICT) [27]. However, the process of obtaining ES-Ag requires harvesting muscle tissue larvae from experimentally infected laboratory animals, which can be resource-intensive and labor-intensive. Recombinant proteins offer a promising alternative to ES antigens, as they can be efficiently produced in large quantities using bacterial expression systems and used as antigens in sensitive, specific, and standardized ELISA tests for trichinellosis serodiagnosis. This approach could enhance the accuracy and convenience of diagnosing the disease. Previous studies [28, 29] have suggested that the serine protease of adult *T. spiralis* larvae plays a role in capsule formation and may help protect newborn larvae in the host bloodstream. These findings imply that serine proteases may be involved in intestinal mucosal invasion and could serve as potential targets for early detection of parasitic invasion, as well as for the development of vaccines against trichinellosis infection.

In our study, larvae were detected in the gut starting at 7 dpi, with their numbers increasing from 12 to 91 larvae by 30 dpi. Transformation into muscle tissue was observed between 21 and 30 dpi, and the number of larvae in the gut significantly decreased from 14 to 1.6 larvae. These findings suggest that signs of the disease can be detected between days 7 and 14 using the recombinant serine protease, enabling earlier diagnosis.

These findings are consistent with studies by O.N. Andreyanov, who demonstrated that at a dose of 100 *T. spiralis* larvae, survival in the intestine persists for 18-24 days [30]. Similarly, Pereverzeva et al. observed no intestinal, sexually mature *Trichinella* in mice after day 21 of infection. In their control group, the survival rates of intestinal *Trichinella* were: 54% after 4 days, 44% after 7 days, 21-30% after 15 days, 16.5% after 21 days, and 4-5% between 28-30 days [31], which aligns with our results.

No encapsulated larvae were observed by day 30 post-infection. These findings align with those of Pereverzeva, who reported the presence of semi-formed larvae in muscle tissue by day 21, suggesting ongoing transport from the intestine. Notably, even by day 35 post-infection, encapsulation in *Trichinella spiralis* was not observed in mice [32].

Serine protease can be expressed at high levels in encapsulated species (such as *T. spiralis*, *T. nativa*, *T. britovi*, and *T. nelsoni*) but at lower levels in non-encapsulated species (e.g., *T. pseudospiralis*) [33]. Previous studies have shown that both ES-Ag and S-Ag detect infection between days 8 and 14, with recombinant antigens exhibiting similar sensitivity [33, 34, 36]. However, recombinant antigens offer significantly higher specificity, and the more straightforward and standardized production process compared to ES-Ag and S-Ag makes them a preferable alternative for diagnostic applications.

In the present study, the recombinant rTsp-LE antigen demonstrated its diagnostic potential across three animal species (mice, rabbits, and pigs), with average antibody titers of 1:570, 1:2260, and 1:800, respectively. In comparison, when ES-Ag and S-Ag were used, the mean titers were much higher: 1:20480 for ES-Ag and S-Ag in rabbits, and 1:6400 and 1:3940 for pigs, respectively. These findings suggest that the recombinant 11 kDa *T. spiralis* antigen could serve as a viable alternative to ES-Ag and S-Ag for diagnosing trichinellosis. Additionally, no significant difference in antibody detection rates was observed when sera from mice infected with *T. nativa* were tested using the recombinant antigen, and there was no cross-reactivity with sera from *Echinococcus*-infected animals.

According to the literature, antibodies are typically detectable 4-6 weeks after humans are infected from wild animals. A diagnostically significant result is a fourfold or greater increase in antibody titers in paired sera collected 14-20 days apart. Antibody levels peak between the 2nd and 4th months post-infection and then gradually decline, but can persist for up to 10 years or longer [34].

Garkavi et al. reported that in experimentally infected mice, total antibody concentrations peak at 6 weeks and again at 2-3 months, with IgM peaking on days 30 and 90, and IgA peaking on days 90 and 120 post-infection [35].

The recombinant serine protease protein has recently been actively studied as a component in vaccination. Mice vaccinated with the purified rTspSP-1.3 protein [36] showed an average 39% reduction in muscle larval load compared to the control group.

Recombinant rTsSP protein combined with cholera toxin subunit B (CTB) [37] induced a significant intestinal local sIgA response and a systemic *TsSP*-specific antibody response in vaccinated mice. Moreover, an increased number of goblet cells, acid mucins, and IgA-secreting cells were observed in the small intestine of vaccinated mice. Immune serum against rTsSP specifically recognized the cuticle of various stages of the worm, including muscle larvae, intestinal infective larvae, and adult worms. Vaccinated mice showed a 71.10% reduction in adult worm numbers at 9 dpi and a 62.10% reduction in muscle larvae at 42 dpi. Furthermore, rTsSP vaccination also inhibited the development of intestinal *T. spiralis* and reduced female fecundity.

These findings align with the study by Wang L. et al., who examined the 31 kDa antigen of *T. spiralis* and observed a slightly lower result compared to ES-Ag, with ELISA sensitivity and specificity of 96.67% and 96.87% for the recombinant protein, versus 100% and 98.44% for ES antigens [13]. In a study of four recombinant antigens for *Trichinella*, the r-P1 protein detected antibodies as early as 8 dpi, followed by the r-P2, r-P3, and r-P4 proteins at 10, 14, and 16 dpi, respectively, with antibody levels remaining elevated up to 45 dpi [38]. Similarly, the recombinant serine protease of *T. spiralis* (rTsSP-ZH68) was recognized by sera from infected mice at 8-10 dpi and by sera from early trichinellosis patients at 19 dpi [39].

The recombinant protein rTsp-LE, developed in this study, showed its effectiveness in detecting infection in experimentally infected animals. This antigen demonstrates promise not only for the development of ELISA-based diagnostic tests but also as a potential component for vaccines aimed at reducing the invasive load in animals.

Detection of the intestinal stage of *T. spiralis* on the 7th day post-infection (dpi) allows for the early identification of antibodies, enabling detection of the disease at an early stage. The studied recombinant protein demonstrates high diagnostic value, showing reliable results as early as after the third immunization. The absence of cross-reaction with echinococcosis, along with the ability to detect antibodies to *T. nativa*, supports the potential use of this protein in the development of an ELISA test. Future studies should focus on exploring the use of this protein not only in diagnostic applications but also as a component of a vaccine, or independently, to reduce the parasitic load during infection.

Authors' Contributions

OA: Designed and supervised the study and drafted the manuscript. AG: Statistical analysis and drafted the manuscript. AS, NA: Designed and conducted the study. DhZ: Bioinformatic analysis. FZ: Conducted the study and drafted the manuscript. NG, SYe: Conducted the study and bioinformatic analysis. All authors have read, reviewed, and approved the final manuscript.

Acknowledgements

The study was funded by the Ministry of Science and Higher Education of the Republic of Kazakhstan to frame the Grant Financing for 2024-2026 the project No. AP23489156 "Identification of alternative genes of *Trichinella nativa* immunospecific proteins for latex diagnosticum producing".

Conflicts of Interest

The authors declare that they have no competing interests.

References

- 1 Клинков, АВ, Одолевская, ИМ. (2010). Сравнительный анализ препаратов защитного действия на основе рекомбинантных и экскреторно-секреторных антигенов *Trichinella spp.* При экспериментальном трихинеллезе лабораторных животных. *Российский паразитологический журнал*, 3.
- 2 Murrell, KD, Pozio, E. (2011). Worldwide occurrence and impact of human trichinellosis, 1986-2009. *Emerging Infectious Diseases*, 17(12), 2194-2202. DOI: 10.3201/eid1712.110896.
- 3 European Centre for Disease Prevention and Control. Trichinellosis. In: ECDC. Annual Epidemiological Report for 2020. (2022). Stockholm: ECDC.
- 4 Zhang, XZ, Wang, ZQ, Cui, J. (2022). Epidemiology of trichinellosis in the People's Republic of China during 2009-2020. *Acta Tropica*, 229, 106388. DOI: 10.1016/j.actatropica.2022.106388.
- 5 Year, H., Bory, S., Khieu, V., Caron, Y. (2022). Human trichinellosis in Southeast Asia, 2001-2021. *Food Waterborne Parasitology*, 16, 28. DOI: 10.1016/j.fawpar.2022.e00171.
- 6 Zhang, N., Li, W., Fu, B. (2018). Vaccines against *Trichinella spiralis*: Progress, challenges and future prospects. *Transboundary and Emerging Diseases*, 65(6), 1447-1458. DOI: 10.1111/tbed.12917.
- 7 Centers of Disease control and prevention (CDC) Clinical Overview of Trichinellosis (2024). <https://www.cdc.gov/trichinellosis/hcp/clinical-overview/index.html#:~:text=Diagnosis,or%20specific%20antibody%20in%20serum>.
- 8 Gamble, HR, Patrascu, IV. (1996). Whole blood, serum, and tissue fluids in an enzyme immunoassay for swine trichinellosis. *Journal of Food Protection*, 59(11), 1213-1217. DOI: 10.4315/0362-028X-59.11.1213.
- 9 Gottstein, B., Pozio, E., Nöckler, K. (2009). Epidemiology, diagnosis, treatment, and control of trichinellosis. *Clinical Microbiology Reviews*, 22(1), 127-145. DOI: 10.1128/CMR.00026-08.
- 10 Rawla, P., Sharma, S. (2024). *Trichinella spiralis* Infection. P. Rawla, StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing.
- 11 Wang, ZQ, Shi, YL, Liu, RD, Jiang, P., Guan, YY, Chen, YD, Cui, J. (2017). New insights on serodiagnosis of trichinellosis during window period: early diagnostic antigens from *Trichinella spiralis* intestinal worms. *Infectious Diseases of Poverty*, 6(1), 41. DOI: 10.1186/s40249-017-0252-z.
- 12 Han, LL, Lu, QQ, Li, YL, Zheng, WW, Ren, P., Liu, RD, Cui, J, Wang, ZQ. (2024). Application of a recombinant novel trypsin from *Trichinella spiralis* for serodiagnosis of trichinellosis. *Parasites Vectors*, 17(1), 9. DOI: 10.1186/s13071-023-06067-7.
- 13 Wang, L., Tian, XY, Sun, GG, Liu, RD, Liu, LN, Zhang, X., Jiang, P., Wang, ZQ, Cui, J. (2015). The Use of Recombinant 31 kDa Antigens of *Trichinella spiralis* for Serodiagnosis of Experimental Trichinellosis. *Iranian Journal of Parasitology*, 10(2), 230-237.
- 14 Supcharoengoon, U., Reamtong, O., Dekumyoy, P., Watthanakulpanich, D., Limpanont, Y., Zhiyue, L., Chaimon, S., Martviset, P., Adisakwattana, P. (2022). Evaluation of indirect-ELISA using eluted antigens from *Trichinella spiralis* muscle larvae for diagnosis of swine trichinellosis. *Acta Tropica*, 235, 106644. DOI: 10.1016/j.actatropica.2022.106644.

- 15 Liu, Y., Xu, N., Li, Y., Tang, B., Yang, H., Gao, W., Liu, M., Liu, X., Zhou, Y. (2021). Recombinant cystatin-like protein-based competition ELISA for *Trichinella spiralis* antibody test in multihost sera. *PLoS Neglected Tropical Diseases*, 15(8). DOI: 10.1371/journal.pntd.0009723.
- 16 Jung, D., Teifke, JP, Karger, A., Michael, K., Venz, S., Wittmann, W., Kindermann, K., Nöckler, K., Mundt, E. (2007). Evaluation of baculovirus-derived recombinant 53-kDa protein of *Trichinella spiralis* for detection of Trichinella-specific antibodies in domestic pigs by ELISA. *Parasitology Research*, 100(3), 429-437. DOI: 10.1007/s00436-006-0296-7.
- 17 Yang, C., Liu, H., Li, X., Sui, S., Liu, Y. (2016). Protective effect of recombinant Trichinella 53-kDa protein in sepsis and the effect on macrophages. *Experimental and Therapeutic Medicine*, 12, 965-968. DOI:10.3892/etm.2016.3354.
- 18 Su, XZ, Prestwood, AK, McGraw, RA. (1991). Cloning and expression of complementary DNA encoding an antigen of *Trichinella spiralis*. *Molecular and Biochemical Parasitology*, 45(2), 331-336. DOI: 10.1016/0166-6851(91)90101-b.
- 19 Wang, B., Wang, ZQ, Jin, J., Ren, HJ, Liu, LN, Cui, J. (2013). Cloning, expression and characterization of a *Trichinella spiralis* serine protease gene encoding a 35.5 kDa protein. *Experimental Parasitology*, 134(2), 148-54. DOI: 10.1016/j.exppara.2013.03.004.
- 20 Nagano, I., Wu, Z., Nakada, T., Matsuo, A., Takahashi, Y. (2001). Molecular cloning and characterization of a 21 kDa protein secreted from *Trichinella pseudospiralis*. *Journal of Helminthology*, 75(3), 273-278. DOI:10.1079/JOH200158.
- 21 The Development of Science-based Guidelines for Laboratory Animal Care: Proceedings of the November 2003 International Workshop. (2004). National Academies of Sciences, Engineering, and Medicine. Washington, DC: The National Academies Press. DOI:10.17226/11138.
- 22 Akibekov, OS, Syzdykova, AS, Lider, LA, Zhumalin, AK, Baibolin, ZK, Zhagipar, FS, Akanova, ZZ, Ibzhanova, AA, Gajimuradova, AM. (2022). Hematological, biochemical, and serological parameters of experimentally infected rabbits with *Trichinella nativa* and *Trichinella spiralis* for early identification of trichinellosis. *Veterinary World*, 15(9), 2285-2292. DOI: 10.14202/vetworld.2022.2285-2292.
- 23 Gamble, HR, Alban, L., Hill, D., Pyburn, D., Scandrett, B. (2019). International Commission on Trichinellosis: Recommendations on pre-harvest control of *Trichinella* in food animals. *Food and Waterborne Parasitology*, 15. DOI: 10.1016/j.fawpar. 2019.e00039.
- 24 Tintori, SC, Sloat, SA, Rockman, MV. (2022). Rapid Isolation of Wild Nematodes by Baermann Funnel. *Journal of Visualized Experiments*, 179. DOI: 10.3791/63287.
- 25 Committee of Sanitary and Epidemiological Control of the Ministry of Healthcare of the Republic of Kazakhstan. (2020). Monitoring the Incidence of Parasitic Infections among the Population of the Republic of Kazakhstan. Available from: <https://www.gov.kz/memleket/entities/kkkbtu/press/news/details/262201?lang=ru>. Retrieved on 10-05-2024.
- 26 Сайдулдин, ТС. (1981). Статистическая обработка результатов серологических реакций. *Ветеринария*, 7, 62-66.
- 27 Liu, RD, Qi, X., Sun, GG, Jiang, P., Zhang, X., Wang, LA, Liu, XL, Wang, ZQ, Cui, J. (2016). Proteomic analysis of *Trichinella spiralis* adult worm excretory-secretory proteins recognized by early infection sera. *Veterinary Parasitology*, 231, 43-46. DOI: 10.1016/j.vetpar.2016.10.008.
- 28 Gamble, HR, Pozio, E., Bruschi, F., Nockler, K., Kapel, C.M., Gajadhar, A.A. (2004). International Commission on Trichinellosis: recommendations on the use of serological tests for the detection of Trichinella infection in animals and man. *Parasite*, 11, 3- 13. DOI: 10.1051/parasite/20041113.
- 29 Xu, J., Bai, X., Wang, LB, Shi, HN, Giessen, JWB, Boireau, P., Liu, MY, Liu, XL. (2017). Immune responses in mice vaccinated with a DNA vaccine expressing serine protease-like protein from the new-born larval stage of *Trichinella spiralis*. *Journal of Parasitology*, 144, 712-719. DOI: 10.1017/S0031182016002493.
- 30 Андреев, ОН. (2009). Биологические свойства изолята *Trichinella spiralis* обыкновенной лисицы, пассированного на лабораторных животных. *Теория и практика борьбы с паразитарными болезнями*, 10.
- 31 Переверзева, ЭВ, Озерецковская, Н., Веретенникова, НЛ. (1974). Реакция белой крови у мышей, зараженных различными штаммами трихинелл. *Wiadomosci Parazytologiczne*, 20(1), 67-80.

32 Переверзева, ЭВ. (1975). Особенности кишечной и мышечной фазы трихинеллеза у белых мышей при экспериментальном угнетении иммунитета. *Wiadomosci Parazytologiczne*, 21, 601-619.

33 Xue, Y., Zhang, B., Wang, N., Huang, HB, Quan, Y., Lu, HN, Zhu, ZY, Li, JY, Pan, TX, Tang, Y., Jiang, YL, Shi, CW, Yang, GL, Wang, CF. (2022). Oral Vaccination of Mice With *Trichinella spiralis* Putative Serine Protease and Murine Interleukin-4 DNA Delivered by Invasive *Lactiplantibacillus plantarum* Elicits Protective Immunity. *Frontiers in Microbiology*, 13. DOI: 10.3389/fmicb.2022.859243.

34 Cui, J., Han, Y., Yue, X., Liu, F., Song, YY, Yan, SW, Lei, JJ, Zhang, X., Jiang, P., Wang, ZQ. (2019). Vaccination of mice with a recombinant novel cathepsin B inhibits *Trichinella spiralis* development, reduces the fecundity and worm burden. *Parasites Vectors*, 12(1), 581. DOI: 10.1186/s13071-019-3833-9.

35 Гаркави, БЛ. (2007). Трихинеллез, вызываемый *Trichinella pseudospiralis* (морфология и биология возбудителя, эпизоотология и эпидемиология, диагностика, меры борьбы и профилактика). *Российский паразитологический журнал*, 2, 35-116.

36 Li, X., Yao, P., Pan, AH, Liu, W., Hu, XC, Wu, ZD, Zhou, XW. (2013). An antigenic recombinant serine protease from *Trichinella spiralis* induces protective immunity in BALB/c mice. *Parasitology Research*, 112(9), 3229-3238. DOI: 10.1007/s00436-013-3500-6.

37 Sun, GG., Lei, JJ, Ren, HN, Zhang, Y., Guo, KX, Long, SR, Liu, RD, Jiang, P., Wang, ZQ, Cui, J. (2019). Intranasal immunization with recombinant *Trichinella spiralis* serine protease elicits protective immunity in BALB/c mice. *Experimental Parasitology*, 201, 1-10. DOI: 10.1016/j.exppara.2019.04.006.

38 Zhai, CC, Liu, XL, Bai, X., Jia, ZJ, Chen, SH, Tian, LG, Ai, L., Tang, B., Liu, MY, Wu, XP, Chen, JX. (2021). Bioinformatic Prediction and Production of Four Recombinant Proteins from Different Developmental Stages of *Trichinella spiralis* and Testing of Their Diagnostic Sensitivity in Mice. *Iranian Journal of Parasitology*, 16(1), 122-135. DOI: 10.18502/ijpa.v16i1.5531.

39 Gao, H., Tang, B., Bai, X., Wang, L., Wu, X., Shi, H., Wang, X. Liu, X., Liu, M. (2018). Characterization of an antigenic serine protease in the *Trichinella spiralis* adult. *Experimental Parasitology*, 195, 8-18. DOI: 10.1016/j.exppara.2018.09.009.

References

1 Klinkov, AV, Odoevskaya, IM. (2010). Sravnitel'nyj analiz preparatov zashchitnogo dejstviya na osnove rekombinantnyh i ekskretorno-sekretornyh antigenov *Trichinella spp.* Pri eksperimental'nom trihinelleze laboratornyh zhivotnyh. *Rossijskij parazytologicheskij zhurnal*, 3. [in Russ].

2 Murrell, KD, Pozio, E. (2011). Worldwide occurrence and impact of human trichinellosis, 1986-2009. *Emerging Infectious Diseases*, 17(12), 2194-2202. DOI: 10.3201/eid1712.110896.

3 European Centre for Disease Prevention and Control. Trichinellosis. In: ECDC. Annual Epidemiological Report for 2020. (2022). Stockholm: ECDC.

4 Zhang, XZ, Wang, ZQ, Cui, J. (2022). Epidemiology of trichinellosis in the People's Republic of China during 2009-2020. *Acta Tropica*, 229, 106388. DOI: 10.1016/j.actatropica.2022.106388.

5 Year, H., Bory, S., Khieu, V., Caron, Y. (2022). Human trichinellosis in Southeast Asia, 2001-2021. *Food Waterborne Parasitology*, 16, 28. DOI: 10.1016/j.fawpar.2022.e00171.

6 Zhang, N., Li, W., Fu, B. (2018). Vaccines against *Trichinella spiralis*: Progress, challenges and future prospects. *Transboundary and Emerging Diseases*, 65(6), 1447-1458. DOI: 10.1111/tbed.12917.

7 Centers of Disease control and prevention (CDC). Clinical Overview of Trichinellosis (2024). <https://www.cdc.gov/trichinellosis/hcp/clinical-overview/index.html#:~:text=Diagnosis,or%20specific%20antibody%20in%20serum>.

8 Gamble, HR, Patrascu, IV. (1996). Whole blood, serum, and tissue fluids in an enzyme immunoassay for swine trichinellosis. *Journal of Food Protection*, 59, 1213-1217. DOI: 10.4315/0362-028X-59.11.1213.

9 Gottstein, B., Pozio, E., Nöckler, K. (2009). Epidemiology, diagnosis, treatment, and control of trichinellosis. *Clinical Microbiology Reviews*, 22(1), 127-145. DOI: 10.1128/CMR.00026-08.

- 10 Rawla, P., Sharma, S. (2024). *Trichinella spiralis* Infection. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing.
- 11 Wang, Z, Shi, YL, Liu, RD, Jiang, P., Guan, YY, Chen, YD, Cui, J. (2017). New insights on serodiagnosis of trichinellosis during window period: early diagnostic antigens from *Trichinella spiralis* intestinal worms. *Infectious Diseases of Poverty*, 6(1), 41. DOI: 10.1186/s40249-017-0252-z.
- 12 Han, LL, Lu, QQ, Li, YL, Zheng, WW, Ren, P., Liu, RD, Cui, J., Wang, ZQ. (2024). Application of a recombinant novel trypsin from *Trichinella spiralis* for serodiagnosis of trichinellosis. *Parasites Vectors*, 17, 9. DOI: 10.1186/s13071-023-06067-7.
- 13 Wang, L., Tian, XY, Sun, GG, Liu, RD, Liu, LN, Zhang, X., Jiang, P., Wang, ZQ, Cui, J. (2015). The Use of Recombinant 31 kDa Antigens of *Trichinella spiralis* for Serodiagnosis of Experimental Trichinellosis. *Iranian Journal of Parasitology*, 10(2), 230-237.
- 14 Supcharoengoon, U., Reamtong, O., Dekumyoy, P., Watthanakulpanich, D., Limpanont, Y., Zhiyue, L., Chaimon, S., Martviset, P., Adisakwattana, P. (2022). Evaluation of indirect-ELISA using eluted antigens from *Trichinella spiralis* muscle larvae for diagnosis of swine trichinellosis. *Acta Tropica*, 235, 106644. DOI: 10.1016/j.actatropica.2022.106644.
- 15 Liu, Y., Xu, N., Li, Y., Tang, B., Yang, H., Gao, W., Liu, M., Liu, X., Zhou, Y. (2021). Recombinant cystatin-like protein-based competition ELISA for *Trichinella spiralis* antibody test in multihost sera. *PLoS Neglected Tropical Diseases*, 15(8). DOI: 10.1371/journal.pntd.0009723.
- 16 Jung, D., Teifke, JP, Karger, A., Michael, K., Venz, S., Wittmann, W., Kindermann, K., Nöckler, K., Mundt, E. (2007). Evaluation of baculovirus-derived recombinant 53-kDa protein of *Trichinella spiralis* for detection of *Trichinella*-specific antibodies in domestic pigs by ELISA. *Parasitology Research*, 100(3), 429-437. DOI: 10.1007/s00436-006-0296-7.
- 17 Yang, C., Liu, H., Li, X., Sui, S., Liu, Y. (2016). Protective effect of recombinant *Trichinella* 53-kDa protein in sepsis and the effect on macrophages. *Experimental and Therapeutic Medicine*, 12, 965-968. DOI: 10.3892/etm.2016.3354.
- 18 Su, XZ, Prestwood, AK, McGraw, RA. (1991). Cloning and expression of complementary DNA encoding an antigen of *Trichinella spiralis*. *Molecular and Biochemical Parasitology*, 45(2), 331-336. DOI: 10.1016/0166-6851(91)90101-b.
- 19 Wang, B., Wang, ZQ, Jin, J., Ren, HJ, Liu, LN, Cui, J. (2013). Cloning, expression and characterization of a *Trichinella spiralis* serine protease gene encoding a 35.5 kDa protein. *Experimental Parasitology*, 134(2), 148-154. DOI: 10.1016/j.exppara.2013.03.004.
- 20 Nagano, I., Wu, Z., Nakada, T., Matsuo, A., Takahashi, Y. (2001). Molecular cloning and characterization of a 21 kDa protein secreted from *Trichinella pseudospiralis*. *Journal of Helminthology*, 75(3), 273-278. DOI:10.1079/JOH200158.
- 21 The Development of Science-based Guidelines for Laboratory Animal Care: Proceedings of the November 2003 International Workshop. (2004). *National Academies of Sciences, Engineering, and Medicine*. Washington, DC: The National Academies Press. DOI: doi.org/10.17226/11138.
- 22 Akibekov, OS, Syzdykova, AS, Lider, LA, Zhumalin, AK, Baibolin, ZK, Zhagipar, FS, Akanova, ZZ, Ibzhanova, AA, Gajimuradova, AM. (2022). Hematological, biochemical, and serological parameters of experimentally infected rabbits with *Trichinella nativa* and *Trichinella spiralis* for early identification of trichinellosis. *Veterinary World*, 15(9), 2285-2292. DOI: 10.14202/vetworld.2022.2285-2292.
- 23 Gamble, HR., Alban, L., Hill, D., Pyburn, D., Scandrett, B. (2019). International Commission on Trichinellosis: Recommendations on pre-harvest control of *Trichinella* in food animals. *Food and Waterborne Parasitology*, 15. DOI: 10.1016/j.fawpar. 2019.e00039.
- 24 Tintori, SC, Sloat, SA, Rockman, MV. (2022). Rapid Isolation of Wild Nematodes by Baermann Funnel. *Journal of Visualized Experiments*, 179. DOI: 10.3791/63287.
- 25 Komitet sanitarno-epidemiologicheskogo nadzora Ministerstva zdavoohraneniya Respubliki Kazahstan. Monitoring zaboлеваemosti parazitarnymi infekciyami sredi naseleniya Respubliki Kazahstan. (2020). [in Russ].
- 26 Saiduldin TS. (1981). Statisticheskaya obrabotka rezultatov serologicheskikh reakcii. *Veterinariya*, 7, 62-66.
- 27 Liu, RD, Qi, X., Sun, GG, Jiang, P., Zhang, X., Wang, LA, Liu, XL, Wang, ZQ, Cui, J. (2016). Proteomic analysis of *Trichinella spiralis* adult worm excretory-secretory proteins recognized by early infection sera. *Veterinary Parasitology*, 231, 43-46. DOI: 10.1016/j.vetpar.2016.10.008.

28 Gamble, HR, Pozio, E., Bruschi, F., Nockler, K., Kapel, CM, Gajadhar, AA. (2004). International Commission on Trichinellosis: recommendations on the use of serological tests for the detection of *Trichinella* infection in animals and man. *Parasite*, 11, 3-13. DOI: 10.1051/parasite/20041113.

29 Xu, J., Bai, X., Wang, LB, Shi, HN, Giessen, JWB, Boireau, P., Liu, MY, Liu, XL. (2017). Immune responses in mice vaccinated with a DNA vaccine expressing serine protease-like protein from the new-born larval stage of *Trichinella spiralis*. *Journal of Parasitology*, 144, 712-719. DOI: 10.1017/S0031182016002493.

30 Andreyanov, ON. (2009). Biologicheskie svoystva izolyata *Trichinella spiralis* obyknovlennoj lisicy, passirovannogo na laboratornyh zhivotnyh. *Teoriya i praktika bor'by s parazitarnymi boleznyami*, 10.

31 Pereverzeva, EV, Ozereckovskaya, NN, Veretennikova, NL. (1974). On the developmental particularities of *T.spiralis* larvae isolated from the racoon (procyon lotor) muscles – in albino mice. *Wiadomosci Parazytologiczne*, 20(1), 67-80.

32 Pereverzeva, EV. (1975). Osobennosti kishhechnoj myshechnoi fazy trihinelleza u belyh myshei pri eksperimental'nom ugnetenii immuniteta. *Wiadomosci Parazytologiczne*, 21, 601-619. [In Russ].

33 Xue, Y., Zhang, B., Wang, N., Huang, HB, Quan, Y., Lu, HN, Zhu, ZY, Li, JY, Pan, TX, Tang, Y., Jiang, YL, Shi, CW, Yang, GL, Wang, CF. (2022). Oral Vaccination of Mice With *Trichinella spiralis* Putative Serine Protease and Murine Interleukin-4 DNA Delivered by Invasive *Lactiplantibacillus plantarum* Elicits Protective Immunity. *Frontiers in Microbiology*, 13. DOI: 10.3389/fmicb.2022.859243.

34 Cui, J., Han, Y., Yue, X., Liu, F., Song, YY, Yan, SW, Lei, JJ, Zhang, X., Jiang, P., Wang, ZQ. (2019). Vaccination of mice with a recombinant novel cathepsin B inhibits *Trichinella spiralis* development, reduces the fecundity and worm burden. *Parasites Vectors*, 12(1), 581. DOI: 10.1186/s13071-019-3833-9.

35 Garkavi, BL. (2007). Trihinellez, vyzyvaemyj *Trihinella pseudospiralis* (morfologiya i biologiya vzbuditelya, epizootologiya i epidemiologiya, diagnostika, mery bor'by i profilaktika). *Rossiiskii parazitologicheskii zhurnal*, 2, 35-116. [In Russ].

36 Li, X., Yao, JP, Pan, AH, Liu, W., Hu, XC, Wu, ZD, Zhou, XW. (2013). An antigenic recombinant serine protease from *Trichinella spiralis* induces protective immunity in *BALB/c* mice. *Parasitology Research*, 112, 3229-3238. DOI: 10.1007/s00436-013-3500-6.

37 Sun, GG, Lei, JJ, Ren, HN, Zhang, Y., Guo, KX, Long, SR, Liu, RD., Jiang, P., Wang, ZQ, Cui, J. (2019). Intranasal immunization with recombinant *Trichinella spiralis* serine protease elicits protective immunity in *BALB/c* mice. *Experimental Parasitology*, 201, 1-10. DOI: 10.1016/j.exppara.2019.04.006.

38 Zhai, CC, Liu, XL, Bai, X., Jia, ZJ, Chen, SH, Tian, LG, Ai, L., Tang, B., Liu, MY, Wu, XP, Chen, JX. (2021). Bioinformatic Prediction and Production of Four Recombinant Proteins from Different Developmental Stages of *Trichinella spiralis* and Testing of Their Diagnostic Sensitivity in Mice. *Iranian Journal of Parasitology*, 16(1), 122-135. DOI: 10.18502/ijpa.v16i1.5531.

39 Gao, H., Tang, B., Bai, X., Wang, L., Wu, X., Shi, H., Wang, X., Liu, X., Liu, M. (2018). Characterization of an antigenic serine protease in the *Trichinella spiralis* adult. *Experimental Parasitology*, 195, 8-18. DOI: 10.1016/j.exppara.2018.09.009.

Herald of Science of S.Seifullin Kazakh Agrotechnical Research University: Veterinary Sciences.
– Astana: S. Seifullin Kazakh Agrotechnical Research University, 2024. – № 4 (008). – P. 16-33.
- ISSN 2958-5430, ISSN 2958-5449

doi.org/ 10.51452/kazatuvc.2024.4(008).1800

UDC 637.5.05/.07

Review article

Meat Quality Control in Beef Production

Shynar K. Sabyrbekova , Gulnaz D. Ilgekbayeva 

Department of Biological Safety, Faculty of Veterinary Medicine,
Kazakh National Agrarian Research University, Almaty, Kazakhstan

Corresponding Author: Shynar K. Sabyrbekova: shynara_85_kz@mail.ru

Co-author: (1: GI) gulnaz66@mail.ru

Received: 29-10-2024 **Accepted:** 17-12-2024 **Published:** 30-12-2024

Abstract

The hazard analysis and critical control points (HACCP) system is primarily a risk assessment method focused on identifying and controlling the risk factors classified as the critical control points. Pathogenic microorganisms can enter the meat supply chain at various points along their path. Live animals can harbor various pathogens, while unhygienic conditions during slaughter and processing introduce external contamination risks, and cross-contamination can occur during handling and cooking. Traditionally, pathogen control is concentrated in the processing stage. Despite its effectiveness, approaches like HACCP do not fully satisfy the consumer demands for food safety. A more comprehensive approach targeting pathogen control at every stage of the supply chain that is, “from farm-to-table,” offers broader coverage and greater integration.

Meat is one of the most perishable food products, providing an ideal environment for the growth of a wide range of pathogenic bacteria. Foodborne illnesses result from the consumption of bacteria, toxins, or cells produced by microorganisms present in food. Past outbreaks have been linked to various stages of the meat production process; hence, it is logical to develop a risk assessment model encompassing all stages, from raising food animals and processing carcasses to meat preparation and consumption.

The primary task in meat production is the evaluation of the microbiological quality of raw meat and the sanitary conditions at processing facilities based on the presence of indicator microorganisms in the meat, equipment, and processing plants. Accordingly, hygienic measures in meat production, processing, and retail aim to ensure meat safety, prevent rapid spoilage, and preserve its quality. HACCP, which heavily relies on prerequisite programs like good hygienic practice and sanitation standard operating procedures, provides enhanced hygiene standards essential for producing safe meat and meat products throughout the entire meat supply chain.

Keywords: critical control points; food products; HACCP; hygienic methods; microbiological quality; slaughterhouse.

Introduction

The hazard analysis and critical control points (HACCP) system is founded on three core principles: hazard identification, determination of critical control points, and implementation of risk mitigation procedures. Additionally, it involves the development of a monitoring system for evaluating the effectiveness of control measures. A critical control point is a stage in the production process where specific interventions can reduce the risk of adverse outcomes. Ineffective control methods should be revised accordingly [1].

For more than twenty years, food safety professionals have been so vigorously promoting HACCP food safety that they have actually overemphasized the usefulness of the HACCP concept [2]. The

global use and success of the HACCP system in the food industry has created false expectations that it can be used successfully at all stages of the food supply chain, from farm to table.

The long global evolution and use of HACCP in food processing facilities has provided a tremendous amount of documentation that the HACCP food safety system is very effective in combating identified foodborne hazards. Its ability to ensure food safety far exceeded that of the quality control system it replaced. The widespread success of HACCP has led regulators, policy makers and consumers to increasingly call for this remarkable tool to be utilized more effectively by applying it throughout the food chain from “farm to table.” Simply put, the supply chain consists of seven stages: animal or crop production, slaughter or harvest, raw food production, processed food production, distribution, catering or retail operations, and consumption. The production of processed products is at the center of this supply chain. It is no coincidence that HACCP originated at the food processing stage - from farm to table.

Meat production is increasing worldwide and according to FAO/WHO estimates [3], it will reach 364 million tons by 2023. The growing demand for meat in developing countries is mainly the result of the rapid progress of societies that are becoming increasingly urbanized.

In recent years, there has been growing concern about the presence of pathogenic microorganisms in meat products despite increased hygiene measures for meat and processed meat products [4]. Contamination can occur during processing, through contact with equipment (e.g., grinders, belts, saws), through contact with food handlers (e.g., hands, knives), and through exposure to other environmental sources (e.g., air, water) [5].

Regulators require meat processing plants to implement hazard analysis and critical control point (HACCP) systems for meat production processes to reduce pathogens [6]. However, since it is difficult or impossible to completely eliminate pathogens from raw meat, the goal of HACCP for meat focuses on reducing and preventing microbial growth [7]. The HACCP system is now used in many regulated sectors of the food industry.

Monitoring microorganisms in meat products is an important step in HACCP programs [8] and proper storage of meat products is critical to control contamination of meat with spoilage microorganisms [9]. It has been proposed that HACCP systems in meat processing plants should be based on microbiological data, with the assessment of the number of indicator organisms in meat products at different stages of processing [10].

For raw meat products, safety and quality can be assessed using indicator microorganisms, including total aerobic platelet count (APC), coliform count (CC), and *Escherichia coli* (ECC) [11]. APC provides an estimate of the total bacterial population. Higher APC is generally associated with lower quality and reduced shelf life. The relationship between APC and the concentration of foodborne pathogens in raw meat is unclear. CC and ECC provide an estimate of faecal contamination and poor sanitation during processing. High CC and ECC values generally correlate with higher levels of foodborne pathogens originating from feces [12]. Meat can be further contaminated or cross-contaminated with various pathogenic bacteria after slaughter, such as during chilling, cutting, deboning, and slicing [13].

Thus, all processing conditions are important factors that can affect the microbiological quality. To improve the safety of finished meat products, more information is needed at the stage of contamination of carcasses or meat cuts in meat processing plants [14]. However, data on seasonal variations and meat type in microbiological testing of meat in different meat processing plants are limited. In particular, microbiological assessment of meat in meat processing plants that have implemented the HACCP system has not been carried out.

Quality Control Programs

The most effective strategy for reducing meat contamination and microbial growth is the implementation of quality control programs, such as good manufacturing practice (GMP) and HACCP. These programs involve identifying indicator microorganisms that can predict the presence of pathogenic microorganisms and spoilage-causing bacteria [15]. Notable bacterial foodborne pathogens, including *Salmonella*, *Campylobacter*, and *verocytotoxigenic Escherichia coli* (VTEC), can accumulate in and be excreted from the gastrointestinal tract of farm animals, including cattle. The typical sequence leading to beef-related foodborne illnesses involves pathogen transmission from cattle to humans through direct or indirect fecal contamination, subsequent cross-contamination, and/or proliferation during the production, processing, and consumption of beef and beef products.

The effective control of beef-borne pathogens necessitates a longitudinal, integrated approach based on the “meat chain” concept. This approach employs the GMP/GHP and HACCP principles, with shared responsibility among all participants in the meat supply chain. It also requires the consideration of resource availability, technical capabilities, consumer attitudes and behavior, and cost-effectiveness [16, 17, 18].

Meat as a Favorable Environment for Microorganism Growth

Meat and meat products provide an exceptionally favorable environment for the growth of pathogenic microorganisms [19]. Its perishable nature and nutrient-rich composition enable meat to support the proliferation of a wide range of pathogenic bacteria [20]. It is vulnerable to contamination at multiple stages, from primary production to the final consumption phase (from farm to fork). Contaminated meat is a major source of foodborne diseases and mortality caused by pathogens entering the body upon ingestion [21]. Foodborne illnesses result from the consumption of bacteria, toxins, or cells produced by microorganisms present in contaminated food [22]. Consequently, retail meat sales are frequently linked to foodborne illnesses if infectious doses are ingested during consumption [23, 24]. Epidemiological and microbiological studies identified cross-contamination during distribution and processing, along with subsequent bacterial growth, as significant contributors to foodborne diseases [25, 26, 27].

Meat and meat products are commonly implicated in food poisoning outbreaks. During production, processing, and storage, these products can become contaminated with pathogenic bacteria, such as *Listeria monocytogenes* [28, 29]. Hygiene and quality during meat handling and processing must be continuously monitored to ensure that meat products adhere to international quality and safety standards [30]. Given its rich nutrient profile, meat is a prime environment for the growth of pathogenic bacteria. The microbiological contamination of carcasses predominantly occurs during handling processes, including skinning, evisceration, processing, storage, and distribution in slaughterhouses and retail outlets [31].

In India, dietary habits have significantly shifted because of rapid urbanization and Westernization, leading to increased consumption of non-vegetarian foods. This shift has resulted in a rising meat demand. Concurrently, consumers have become more vigilant regarding meat quality, freshness, and health aspects [32]. The microbiological quality of meat and meat products is critical for public health. Numerous reports have documented foodborne disease outbreaks linked to meat consumption [33, 34, 35]. Fecal contamination, either through direct deposition or indirect contact via contaminated equipment, workers, facilities, and air, is a primary source of contamination [36]. Pathogens can contaminate meat at various stages of the slaughtering process [37, 38], necessitating appropriate control measures for eliminating or preventing microbial contamination. Among the pathogenic bacteria that can contaminate beef, *Salmonella spp.*, *Listeria monocytogenes*, and *E. coli O157* are frequently associated with foodborne illnesses [39, 40]. It is now widely acknowledged that traditional meat inspection procedures cannot fully ensure that consumers are not exposed to infectious doses of meat-borne pathogens [41]. Therefore, meat inspection authorities worldwide are promoting or mandating the implementation of HACCP systems for meat production processes. While comprehensive HACCP systems should address physical, chemical, and microbiological hazards, the latter remains as a primary concern in meat production [42].

Critical Control Points

In the pre-harvest stage (on the farm), the global recycling of microbial pathogens can be mitigated through the strategic management of agricultural lands and livestock by-products. The fecal shedding of pathogens by farm animals is reduced by employing antimicrobial feed treatments, sourcing new animals from controlled environments, implementing biosecurity measures, optimizing animal welfare (including stress management), maintaining hygienic housing conditions, and utilizing prebiotics, probiotics, competitive exclusion strategies, and vaccination programs.

During the harvest stage (at the slaughterhouse), minimizing transportation time and ensuring optimal pre-slaughter conditions for animals can help reduce cross-contamination via transport vehicles or holding pens. The key measures include maintaining hygienic practices during slaughter and carcass processing, implementing effective sanitation procedures, and potentially applying treatments to hides and/or carcasses to prevent contamination.

In the post-harvest stage (processing, storage, distribution, and consumption), pathogen control in meat products often relies on bactericidal steps, such as cooking, or, where such steps are not employed, on the “multiple hurdles” concept of microbial control. Additional post-harvest risk reduction measures include effective cleaning and sanitation of all facilities involved in meat-processing, maintaining the cold chain, preventing cross-contamination during further processing or cooking, and educating consumers on food hygiene. The effectiveness of control measures varies depending on the pathogen and the stage of the meat supply chain. For instance, some pathogens in beef (e.g., *VTEC*, *Salmonella*) are most effectively controlled through primary production measures combined with enhanced slaughter hygiene. In contrast, other pathogens (e.g., *L. monocytogenes*) are best controlled during the processing and storage stages [43].

Quality monitoring systems are commonly implemented in the food industry to systematically assess indicators at specific points along the technological line and during meat handling [44]. In these systems, operations that significantly affect the microbiological quality of the product must be identified as critical control points (CCPs) and/or quality control points and managed to minimize contamination by pathogenic or spoilage bacteria [45]. Cleaning and disinfection are essential components of biosecurity on livestock farms, reducing animal exposure to foodborne pathogens [46, 47].

Management practices in feedlots can affect animal health, carcass quality, and, potentially, food safety. Feedlots, where cattle from various sources are mixed and housed at high densities, can be sources of contamination by *EHEC O157* (*enterohemorrhagic E. coli O157*) and *Salmonella* spp. Advanced enrichment and isolation techniques demonstrated that the *EHEC O157* prevalence among cattle is significantly higher than that previously estimated, reaching 5.8% [48, 49]. Microbiological testing provides a means of assessing how effectively operators control slaughtering, processing, and production processes to minimize and control contamination [50]. The effectiveness of these systems has been demonstrated. Based on these indicators, corrective actions can also be taken to prevent or reduce potential contamination.

The safety and quality of meat products can be evaluated using various microbial indicators, such as the aerobic mesophilic count and coliforms [51]. The aerobic mesophilic count assesses the general microbial population in the environment and processing equipment, where high contamination levels indicate poor hygienic conditions [52]. Coliforms serve as indicators of inadequate sanitary conditions during food processing, production, and storage, with *E. coli* being a classic indicator of the potential presence of intestinal pathogens and a good measure of the sanitary quality of processed foods [53].

The *Salmonella* spp. transmission from animals to food is a significant concern for human health. It can only be controlled through rigorous cleanliness and adherence to hygienic procedures during slaughter and processing [54]. As natural reservoirs of pathogenic strains, cattle often contaminate ground beef with *EHEC O157* isolated from live animals, carcasses, and retail meat [55]. One study found that cattle housed in feedlots for less than 20 days were 3.4 times more likely to test positive for *EHEC O157* compared to those housed for longer periods [56]. Although *E. coli* is a common and usually harmless inhabitant of the gastrointestinal tract, some strains produce verotoxin that causes severe illness, including diarrhea, hemorrhagic colitis, and potentially fatal kidney failure [57, 58]. Stress responses in cattle, which are potentially linked to feed withdrawal and transportation, may exacerbate these issues. Additionally, *EHEC O157* has been detected in water bodies on many farms, suggesting that water sources may act as long-term reservoirs for this pathogen [59].

“Farm-to-Table” Concept

Meat is susceptible to contamination from both internal and external sources to the animal during slaughter and sale. Live animals can harbor a range of microorganisms on the surfaces that come into contact with their environment. The major contamination sources include animal hides and feces. Additional contamination sources can be the slaughter area, slaughterhouse environment, retail floors, air, and vehicles used for transporting meat from the slaughterhouse [60, 61]. Retail points also contribute to meat contamination [62]. In retail environments, tools like knives, wooden boards, and scales are common sources of bacterial contamination, particularly from *Staphylococcus aureus* and *Shigella* species [63]. Cross-contamination during transport not only affects cattle, but is also a significant source of pathogen transmission in poultry. Despite cleaning and sanitizing efforts, the crates and containers used for poultry transport often remain contaminated with *Salmonella* [64].

Considering that past disease outbreaks are linked to multiple production stages, it is advisable to create a thorough risk assessment model that addresses every stage, from animal rearing and carcass processing to meat preparation and consumption. In pork production, the supply chain encompasses the following stages: pre-harvest (on the farm, during transport, and storage), slaughter (skinning, evisceration, chilling, and production), processing (grinding, treatment, transport, storage, and distribution), consumption (cooking and eating), dose–response (exposure assessment, dose–response function, morbidity, and mortality), and human disease costs [65, 66].

The primary sources of the microbiological contamination of beef carcasses during slaughter include the leakage of intestinal contents and the cross-contamination from hides of slaughtered animals [67, 68]. Fecal contamination is a significant source of microorganisms [69]. The microbiological testing of carcasses is routinely used to verify the HACCP compliance in slaughterhouses. In the European Union (EU), this involves measuring aerobic colony counts and *Enterobacteriaceae* counts known as hygiene indicator organisms to ensure they are within established permissible limits [70].

Cutting and boning operations at meat-processing facilities involve extensive handling, which increases microbial risk due to the following reasons: (a) cross-contamination from hands and tools (e.g., knives, saws, and conveyors); and (b) the transfer of bacteria from the meat surface to its internal parts [71]. However, satisfactory results from slaughter line testing do not guarantee the meat product safety. Additionally, microbial contamination on surfaces in contact with food can sometimes exceed that on carcasses, leading to the accumulation of contamination during the cutting and processing stages [72, 73].

Slaughterhouses are a critical sector in the food industry in terms of potential foodborne illnesses and health hazards, particularly if the food hygiene principles are not strictly followed [74]. The hygiene practices of slaughterhouse workers play a crucial role in contamination. Dirty hands, clothing, and equipment used in carcass handling are contamination sources [75]. A study indicated that 48.4% of slaughterhouse workers did not cover their hair; 29% did not use aprons; and 64.5% wore jewelry (e.g., rings, bracelets, and watches) while working. These findings are consistent with data showing that 61.6% of workers did not cover their hair, and jewelry usage was not regulated [76].

The contamination in slaughterhouses and retail meat shops arises from the use of contaminated water, unhygienic practices (e.g., improper handling), and the use of contaminated surfaces and tools during meat-cutting operations [77].

Transportation can be a stressor for farm animals associated with the shedding of pathogenic bacteria like *EHEC O157* and *Salmonella* spp. in feces, leading to a contamination of trailer floors and bedding [78]. If trailers and bedding are not cleaned and disinfected between trips, they can spread contamination to other farms, the slaughterhouse environment, and other animals [79]. Hygiene rules apply not only during animal housing, but also during transportation. Cattle transported over long distances (more than 160.9 km) are twice as likely to test positive for *E. coli O157* at slaughter compared to those transported at shorter distances [80]. Ineffective vehicle cleaning and sanitizing contributed to contamination in 84% of cattle hides with *E. coli O157*, which had not been detected in any animal at the farm of origin [81].

Hygienic Methods and Quality Control of Meat and Meat Products

As widely recognized, the most significant threats to fresh meat products come from pathogenic bacteria capable of causing illness in humans, including *Salmonella*, *S. aureus*, *L. monocytogenes*, *Campylobacter*, and *E. coli O157*. Particularly, *E. coli O157* can cause food poisoning with only a small number of cells. The primary contamination sources are the slaughter animals themselves, workers, and the working environment. Air contamination through aerosols and water used in carcass processing plays a relatively minor role [82, 83]. Contaminants largely originate from an animal's skin and include microorganisms from the stomach and the intestines, which are released with feces [84]. Foodborne pathogens are a significant public health and economic concern in developed countries [85, 86].

A contamination level of up to 10^5 CFU/cm² indicates good hygienic conditions during slaughter, while higher levels suggest unsatisfactory conditions. Meat contaminated at 10^6 CFU/cm² indicates spoilage, characterized by unpleasant odor and reduced shelf life, while contamination reaching 10^7 CFU/cm² is associated with a noticeable slime formation [30]. Mold counts are used as the indicators of sanitary quality in food production facilities because molds can rapidly proliferate on food residues

adhering to surfaces, consequently posing a potential contamination risk. Some molds produce toxic metabolites known as mycotoxins, which can pose health risks to humans and animals [87].

Hygienic methods and quality control measures for meat and meat products are recommended in many countries, particularly in the food service sector [88]. Without proper hygiene control, slaughterhouse and meat-processing environments can become significant sources of bacterial contamination [89].

The primary objective in meat production is the assessment of the microbiological quality of raw meat and the sanitary conditions at processing facilities based on the presence of indicator microorganisms in the meat, equipment, and processing plants. Identifying the key contamination points and introducing microorganisms into meat allow specialists responsible for production hygiene to implement proper manufacturing practices at meat-processing facilities, thereby improving a product's microbiological quality [90]. Microbiological carcass testing is commonly used to verify the HACCP compliance in slaughterhouses. In the EU, this involves measuring the aerobic colony and *Enterobacteriaceae* counts, known as hygiene indicator organisms, to ensure they are within specified permissible limits [91].

Hygienic measures in meat production, processing, and retail aim to ensure meat safety, prevent rapid spoilage, and maintain quality. HACCP, which heavily relies on prerequisite programs, such as GHP and sanitation standard operating procedures, ensures improved production hygiene conditions necessary for safe meat and meat products throughout the meat supply chain. By the end of the 20th century, HACCP had been mandated, implemented, and continuously used by every meat company involved in international trade [92].

Under EU legislation that is, regulation EC nos. 852/2004, 853/2004, and 854/2004, food business operators at all stages of the production chain bear the primary legal responsibility for ensuring the safety of the food they produce. While HACCP-based procedures are not required at the primary production stage (on the farm), farmers are expected to produce food with hygiene considerations in mind using an approach similar to HACCP. Although the classic HACCP approach is not fully applicable to food safety control on meat farms, the HACCP principles can still be applied to identify hazards and potential corrective actions. Food safety systems for beef should include at least: (1) full animal traceability; (2) provision of clean, uncontaminated feed; (3) provision of clean, uncontaminated water; (4) hygienic animal housing/living conditions; (5) clean livestock; (6) animal health, welfare, and disease prevention; (7) proper waste disposal; (8) hygienic transport and handling; and (9) biosecurity [93, 94].

Foodborne diseases are more common in developing countries because of the widespread improper food handling and sanitation practices, inadequate food safety laws, weak regulatory systems, lack of financial resources for safer equipment, and insufficient education for food industry workers [95].

The bacterial composition of meat serves as an acceptable indicator of its hygienic quality. The poor infrastructure at slaughterhouses, unsanitary animal handling, and improper carcass handling contribute to high bacterial loads in meat. Therefore, evaluating bacterial counts can indicate potential threats to human health [96].

Conclusion

A higher meat contamination level by these microorganisms reflects the poor hygiene and sanitation practices applied at the slaughterhouse during transportation to meat-processing plants and during processing at meat plants [97].

However, the lack of specific critical control points that can eliminate or control the identified hazards hinders the effective use of HACCP throughout the supply chain. Food safety measures can be used at every stage of the supply chain, but most of these measures will be pre-programs rather than critical control points from the HACCP system [98]. The most pressing food safety issues in the food industry today are caused by the presence of *Escherichia coli* O157:H7 and *Salmonella* in raw meat and poultry products and in processed foods. Attempts to control these pathogens either at the “farm” or “table” level of the supply chain have failed due to the lack of effective control measures at these stages available at the food processing stage.

Cross-contamination and other poor food handling practices can also lead to foodborne illness. Thus, the idea that HACCP can be effectively applied from farm to fork has been reconsidered.

The microbiological quality of beef and meat products is highly dependent on the hygienic conditions prevailing during their production and processing. Without proper hygienic control, the environment in slaughterhouses and butcher shops can be an important source of microbiological contamination.

The impact of HACCP on process hygiene performance in slaughterhouses has been previously studied [99, 100]

Specific studies were conducted to determine whether mandatory implementation of HACCP systems in meat processing and retail establishments has any significant impact on their microbiological indicators of process hygiene [101]. The reduction in bacterial counts on food contact surfaces, butchers' hands and refrigeration units was found to provide strong evidence of improved process hygiene performance and justifies the adoption of GHP and SSOP in meat plants. This relatively large-scale study showed that the implementation of HACCP was associated with improved process hygiene in meat plants. However, this finding is necessarily limited in scope as it is not possible at this time to determine whether such improvements would have a similar positive effect on the incidence of meat borne diseases. Further research is clearly needed to establish this.

This information on the key microbiological contamination points in the beef processing chain will assist hygiene professionals in such establishments in developing appropriate hygiene procedures to prevent or reduce microbiological contamination of beef and meat products.

Authors' Contributions

GI: Conceptualized and designed the study, analyzed the data, and drafted the manuscript. SS: Conducted an extensive literature review and analyzed the data. Both authors reviewed and approved the final manuscript.

Acknowledgments

These studies were conducted on the initiative topic registered in the National Center of State Scientific and Technical Expertise of the Republic of Kazakhstan "Determination of CCPs in beef production in the conditions of the south and southeast of the Republic of Kazakhstan" (state registration number 0124RKI0140).

References

- 1 Villarroel, A., Dargatz, DA, Lane, VM, McCluskey, BJ, Salman, MD. (2007). Suggested outline of potential critical control points for biosecurity and biocontainment on large dairy farms. *Food Control*, 203(6), 663-669. DOI: 10.2460/javma.230.6.808.
- 2 Sperber, WH. (2005). HACCP does not work from Farm to Table. *Food Control*, 16, 6, 511-514. DOI: 10.1016/j.foodcont.2003.10.013
- 3 Joint FAO/WHO Food Standards Program Codex Alimentarius Commission, Forty-sixth Session. FAO headquarters, Rome, Italy, 27 November to 2 December 2023.
- 4 Bae, YY, Choi, YM, Kim, MJ, Kim, KH, Kim, BC, Rhee, MS. (2011). Application of supercritical carbon dioxide for microorganism reductions in fresh pork. *J. Food Saf.*, 31, 511-517. DOI: 10.1111/j.1745-4565.2011.00328.x.
- 5 Kim, JH, Yim, DG. (2016). Assessment of the microbial level for livestock products in retail meat shops implementing HACCP system. *Korean J Food Sci An.*, 36, 594-600. DOI: 10.5851/kosfa.2016.36.5.594.
- 6 Agência Nacional de Vigilância Sanitária (2019). Instrução Normativa nº60 de 23 de dezembro de 2019, Estabelece as listas de padrões microbiológicos para alimentos. *Diário Oficial República Federativa do Brasil*, 249(1), 133.
- 7 Ribeiro Júnior, JC., Dias, BP, Nascimento, Nascimento, AL, Cabral Silva, JP, Cordeiro, RF, Oliveira Lobo, CM. (2024). Effects of washing sanitation standard operating procedures on the microbiological quality and safety of cattle carcasses. *Food Control*, 166. DOI: 10.1016/j.foodcont.2024.110745.
- 8 Clinquart, A., Ellies-Oury, P., Hocquette, F., Guillier, L., Santé-Lhoutellier, V., Prache, S. (2022). Review: On-farm and processing factors affecting bovine carcass and meat quality. *Animal*, 1, 100426. DOI: 10.1016/j.animal.2021.100426.
- 9 Kim, JH, Yim, DG. (2017). Microbial levels for food contact and environmental surfaces in meat processing plants and retail shops. *Food Sci Biotechnol.*, 26, 299-302.

10 Ahmed, AH, Al-Mahmood, OA. (2023) Food Safety Programs that should be Implemented in Slaughterhouses: Review. *Journal of Applied Veterinary Sciences*, 8(2), 80-88. DOI: 10.21608/JAVS.2023.185918.1208.

11 Wang, X., Bouzembrak, Y., Oude Lansink, AGJM, van der Fels-Klerx, HJ. (2021). Application of machine learning to the monitoring and prediction of food safety: A review. *Comprehensive Reviews in Food Science and Food Safety*, 21: 1, 416-434. DOI: 10.1111/1541-4337.

12 Abou ElNour-Mona M, Sakr-Ebtehag, AE. (2020). Isolation and Molecular. Identification of Some Food Borne Pathogens from Raw and Processed Meats: In vitro Synergistic Probability of Lactobacillus CFS and Antibiotic. *Egypt. J.Bot.*, 60(2), 405-421. DOI: 10.21608/EJBO.2019.19837.1395.

13 Kuzuoka, K., Kawai, K., Yamauchi, S., Okada, A., Inoshima, Y. (2020). Chilling control of beef and pork carcasses in a slaughterhouse based on causality analysis by graphical modelling. *Food Control*, 32, 106075. DOI: 10.1016/j.foodcont.2020.107353.

14 Prache, S., Schreurs, N., Guillier, L. (2022). Review: Factors affecting sheep carcass and meat quality attributes. *Animal*, 1:100330. DOI: 10.1016/j.animal.2021.100330.

15 Mansur, AR, Song, EJ, Cho, YS, Nam, YD, Choi, YS, Kim, DO. (2019). Comparative evaluation of spoilage-related bacterial diversity and metabolite profiles in chilled beef stored under air and vacuum packaging. *Food Microbiol*, 77, 166-72.

16 *European Surveillance of Veterinary Antimicrobial Consumption (ESVAC)*. (5.02.2023). <https://www.ema.europa.eu/en/veterinary-regulatory/overview/antimicrobial-resistance/european-surveillance-veterinary-antimicrobial-consumption-esvac>.

17 Abd El-Tawab, A., Maarouf, AA, El-Hofy-Fatma, I., Ahmed-Nesma, MG. (2020). Phenotypic characterization of some food poisoning bacteria isolated from meat and meat products in Kaliobia, Egypt. *Benha Vet. Med. J.*, 38(2), 146-151. DOI: 10.21608/bvmj.2020.25678.1186.

18 Røtterud, OJ, Gravning, GE, Hauge, SJ, Alvseike, O. (2020). Hygiene performancerating-an auditing scheme for evaluation of slaughter hygiene. *Methods X*, 20:7, 100829. DOI: 10.1016/j.mex.2020.100829.

19 Korany, SH, Tolba, KS, Sobhy, HM, Hekal, SA. (2021). Assuring the Safety of Local and Imported Beef Meat from Different Slaughterhouses in Egypt. *Advances in Animal and Veterinary Sciences*, 9(9). DOI: 10.17582/journal.aavs/2021/9.9.1472.1482.

20 Birhanu, W., Weldegebriel, S., Bassazin, G., Mitku, F., Birku, L., Tadesse, M. (2017). Assesment of microbiological quality and meat handling practices in butcher shops and abattoir found in Gondar town, Ethiopia. *International Journal of Microbiological Research*, 8:2, 59-68.

21 Food Control 140 (2022). 109143 8 FAO & WHO. (2020). COVID-19 and food safety: Guidance for food businesses. https://a pps.who.int/iris/bitstream/handle/10665/331705/WHO-2019-nCoV-Food_Safety-2020.

22 Sagawa, R., Rodrigues, YM, Nascimento, CA, Ribeiro, J., Oliveira, MS, Muller, AC, Conti, ACM, Ribeiro Júnior, JC. (2021). Impact of the pre-slaughter period on the contamination of bovine leather and the operational sanitary procedure for skinning on the quality and microbiological safety of the carcass. *Semina: Ciênc. Agrár.*, 43, 4. DOI: 10.5433/1679-0359.2022v43n4p1835

23 Cavalcanti, AA, Limeira, CH, Siqueira, IN, de Lima, AC, de Medeiros, FJP, de Souza, JG, de Araújo Medeiros, NG, Abrahão Alves de, OF, de Melo, MA. (2022). The prevalence of *Listeria monocytogenes* in meat products in Brazil: A systematic literature review and meta-analysis. *Research in Veterinary Science*, 145, 169-176.

24 Perez-Rodriguez, F., Castro, R., Posada-Izquierdo, GD, Valero, A., Carrasco, E., Garcia-Gimeno, RM, Zurera, G. (2010). Evaluation of hygiene practices and microbiological quality of cooked meat products during slicing and handling at retail. *MeatSci.*, 86, 479-485. DOI: 10.1016/j.meatsci.2010.05.0387.

25 Alfian, G., Syafrudin M., Farooq, U., Ma'arif, MR, Syaekhoni, MA, Fitriyani, NL, Lee, J., Rhee, J. (2020). Improving efficiency of RFID-based traceability system for perishable food by utilizing IoT sensors and machine learning model. *Food Control*, 110, 107016. DOI: 10.1016/j.foodcont.2019.107016.

26 Cevallos-Almeida, M., Burgos-Mayorga, A., Gómez, CA, Lema-Hurtado, JL, Lema, L., Calvache, I., Jaramillo, C., Ruilova, IC, Martínez, EP, Estupiñán, P. (2021). Association between animal welfare indicators and microbiological quality of beef carcasses, including *Salmonella* spp., from a slaughterhouse in Ecuador. *Veterinary World*, 14(4), 918-925. DOI: 10.14202/vetworld.2021.918-925.

- 27 Oliveira, MS, Santos, IGC, Dias, BP, Nascimento, CA, Rodrigues, ÉM, Ribeiro, JC, Jr., Alfieri, AA, Alexandrino, B. (2021). Hygienic-health quality and microbiological hazard of clandestine Minas Frescal cheese commercialized in north Tocantins, Brazil. *Semina: Ciências Agrárias*, 42(2), 679-694. DOI: 10.5433/1679-0359.2021v42n2p679.
- 28 Birhanu, S., Menda, S. (2017). Hygienic Handling and Processing of Raw Beef Meat at Slaughter Houses and Meat Stalls in Gojjam Area. (*IJVHRS*), 5(8): 213-216. <https://doi.org/10.19070/2332-2748-1700042>.
- 29 Antic, D., Houf, K., Michalopoulou, E., Blagojevic, B. (2021). Beef abattoir interventions in a risk-based meat safety assurance system. *Meat Science*, 182, 108622. DOI: 10.1016/j.meatsci.2021.108622.
- 30 Nakamura, A., Takahashi, H., Kondo, A., Koike, F., Kuda, T., Kimura, B., Kobayashi, M. (2022). Distribution of psychrophilic microorganisms in a beef slaughterhouse in Japan after cleaning. *PLoS ONE*, 17(8), e0268411. DOI: 10.1371/journal.pone.0268411.
- 31 Antic, D., Houf, K., Michalopoulou, E., Blagojevic, B. (2021). Beef abattoir interventions in a risk-based meat safety assurance system. *Meat Science*, 182, 108622. DOI: 10.1016/j.meatsci.2021.108622.
- 32 Gutema, FD, Abdi, RD, Agga, GE, Firew, S., Rasschaert, G., Mattheus, W., Crombe, F., Duchateau, L., Gabriël, S., De Zutter, L. (2021). Assessment of beef carcass contamination with Salmonella and E. coli O 157 in slaughterhouses in Bishoftu, Ethiopia. *International Journal of Food Contamination*, 8(1), 1-9. DOI: 10.1186/s40550-021-00082-1.
- 33 De Filippis, F, Storia, AL, Villani, F, Ercolini, D. (2019). Strain-level diversity analysis of *Pseudomonas fragi* after in situ pangenome reconstruction shows distinctive spoilage associated metabolic traits clearly selected by different storage conditions. *Appl Environ Microbiol*, 85(1).
- 34 Carrasco-García, AA, Pardío-Sedas, VT, León-Banda, GG, Ahuja-Aguirre, C., Paredes-Ramos, P., Hernández-Cruz, BC, Murillo, VV. (2020). Effect of stress during slaughter on carcass characteristics and meat quality in tropical beef cattle. *Asian-Australasian Journal of Animal Sciences*, 33(10), 1656-1665. DOI: 10.5713/ajas.19.0804.
- 35 Nakamura, A., Takahashi, H., Koike, F., Kuda, T., Kobayashi, M. (2023). Transition of microbial contamination on the surface of carcass during the cattle slaughter process., *Food Microbiology*, 104245.
- 36 Calle, A., Carrascal, AK, Patiño, C., Carpio, C., Echeverry, A., Brashears, M. (2021). Seasonal effect on Salmonella, Shiga toxin-producing E. coli O157: H7 and non-O157 in the beef industry in Colombia, South America. *Heliyon*, 7(7), e07547. DOI: 10.1016/j.heliyon.2021.
- 37 Gill, CO. (2007). Microbiological conditions of meats from large game animals and birds. *Meat Science*, 77, 149-160.
- 38 Durmuşoğlu, H., İncili, GK, Demir, P, İlhak, Oİ. (2020). Effects of workers' hand washing and knife disinfection practices on microbiological quality of small animal carcasses in slaughterhouse environment. *Food Process Preserv.* 44(12), 1-7.
- 39 Kuzuoka, K., Kawai, K., Yamauchi, S., Okada, A., Inoshima, Y. (2018). Chilling control of beef and pork carcasses in a slaughterhouse based on causality analysis by graphical modelling. *Food Control*, 118, 107353.
- 40 Nakamura, A., Takahashi, H., Otomo, K., Mizuno, Y., Kuda, T., Kimura, B., Koike, F., Kobayashi, M. (2021). Dynamics of microbiota in Japanese Black beef stored for a long time under chilled conditions. *Food Microbiol*, 100, 103849.
- 41 Akaike, H., Nagai, M., Okatani, AT, Morita, Y. (2023). Food safety, livestock health, and productivity of a farm fattening Japanese black cattle following implementation of a certificated Hazard Analysis and Critical Control Point (HACCP) system. *J Vet Med Sci.*, 7:85(9), 985-989.
- 42 Alvseike, O., Røssvoll, E., Røtterud, OJ, Nesbakken, T., Skjerve, E., Prieto, M., Sandberg, M., Johannessen, G., Økland, M., Urdahl, AM, Hauge, SJ. (2019). Slaughter hygiene in European cattle and sheep abattoirs assessed by microbiological testing and Hygiene Performance Rating. *Food Control*, 101, 233-240.
- 43 Mallhi, IY, Sohaib, M., Khan, AU, Nawaz, M. (2019). Evaluating food safety knowledge, practices, and microbial profile of meat in abattoirs and butchery shops in Lahore, Pakistan. *Journal of Food Safety*, 39(5).
- 44 Odeyemi, OA, Alegbeleye, OO, Strateva, M, Stratev, D. (2020). Understanding spoilage microbial community and spoilage mechanisms in foods of animal origin. *Compr Rev Food Sci Food Saf.*, 19(2), 311-31.

- 45 Webb, NB, Marsden, JL. (1995). *Relationship of the HACCP system to Total Quality Management*. In: Person, AM, Dutson, TR. (Eds.), *Blackie Academic*, London, 10, 156-181.
- 46 Blagojević, B., Nesbakken, T., Alvseike, O., Vågsholm, I., Antic, D., Johlerf, S., Houf, K., Meemken, D., Nastasijevic, I., Vieira Pinto, M., Antunowic, B., Georgiev, M., Alban, L. (2021). Challenges and opportunities in the implementation of new meat inspection systems in Europe. *Trends in Food Science & Technology*, 116, 460-467.
- 47 Panea, B., Ripoll, G. (2020). Designing food safety standards in beef jerky production process with the application of hazard analysis critical control point (HACCP). *Nutrition & Food Science*, 50, 333-347.
- 48 Elder, RO, Keen, JE, Siragusa, GR, Barkocy-Gallagher, GA, Koohmaraie, M., Laegreid, WW. (2000). Correlation of Enterohemorrhagic *Escherichia coli* O157 prevalence in feces, hides, and carcasses of beef cattle during processing. *Proc. Natl. Acad. Sci.*, 97(7), 2999-3003.
- 49 Antic, D., Houf, K., Michalopoulou, E., Blagojevic, B. (2000). Beef abattoir interventions in a risk-based meat safety assurance system. *Meat Sci.*, 182, 108622. DOI: 10.1016/j.meatsci.2021.108622.
- 50 MIG, Microbiological Criteria Information for the UK Meat Industries, MIG, Moscow, Russia, 2006.
- 51 Ghafir, Y., China, B., Dierick, K., Zutter, LD, Daube, G. (2008). Hygiene indicator microorganisms for selected pathogens on beef, pork, and poultry meats in Belgium. *J. Food Prot.* 71(1), 35-45.
- 52 Gill, CO, Deslandes, B., Rahn, K., Houde, A., Bryant, J. (1998). Evaluation of the hygienic performances of the processes for beef carcass dressing at 10 packing plants. *J. Appl. Microbiol.* 84(6), 1050-1058.
- 53 Eisel, WG, Linton, RH, Muriana, PM. (1997). A survey of microbial levels for incoming raw beef, environmental sources, and ground beef in a red meat processing plant. *Food Microbiol.* 14:3, 273-282.
- 54 Wray, C., Davies, RH. (2000). *Salmonella in domestic animals*. CABI Publishing, Wallingford, UK. 184.
- 55 Lynn, TV, Hancock, DD, Besser, TE, Harrison, JH, Rice, DH, Stewart, NT, Rowan, LL. (1998). The occurrence and replication of *Escherichia coli* in cattle feeds. *J. Dairy Sci.*, 81(4), 1102-1108.
- 56 U.S. Department of Agriculture. (1997). USDA-APHIS: factors associated with *Escherichia coli* O157 in feces of feedlot cattle. Centers for Epidemiology and Animal Health, Fort Collins, Colo.
- 57 Armstrong, GL, Hollingsworth, J., Morris, JG Jr. (1996). Emerging foodborne pathogens: *Escherichia coli* O157:H7 as a model of entry of a new pathogen into the food supply of the developed world. *Epidemiol. Rev.*, 18(1), 29-51.
- 58 ANZFS (Australia New Zealand Food Standards Code) Microbiological Limits for Food - Schedule 27 - F2021C00605 (2021).
- 59 Laegreid, WW, Elder, RO, Keen, JE. (1999). Prevalence of *Escherichia coli* O157:H7 in range beef calves at weaning. *Epidemiol. Infect.* 123, 291-298.
- 60 Cooper, K. (1999). The Plant Environment Counts: Protect your Product through Environmental, Sampling. *Meat and Poultry*, 5, 5.
- 61 Sofos, JN, Kochevar, SL, Bellinger, GR, Buege, DR, Hancock, DD, Ingham, SC, Morgan, JB, Reagan, JO, Smith, GC. (1999). Sources and extent of microbiological contamination of beef carcasses in seven United States slaughtering plants. *J Food Prot.* 62(2), 140-5.
- 62 Sudhakar, B, Paturkar, Waskar, VS, Zende, RJ. (2009). Bacteriological screening of environmental sources of contamination in an abattoir and the meat shops in Mumbai, India. *As. J. Food Ag-Ind.*, 2(3), 277-287.
- 63 Ali, N., Farooqui, A., Khan, A., Kazmi, S. (2010). Microbial contamination of raw meat and its environment in retail shops in Karachi, Pakistan. *Journal of Infection in Developing Countries*, 4: 6, 382-388.
- 64 Arquias, RK, Seixas, DB. (2021). Riscos do consumo de carne não inspecionada e as principais características sensoriais analisadas pelos consumidores-revisão de literatura. Universitário FG. <https://repositorio.animaeducacao.com.br/handle/ANIMA/13584>.
- 65 Miller, GY, Liu, X., McNamara, PE, Barber, DA. (2005). The influence of *Salmonella* in pigs preharvest and during pork processing on human health costs and risks from pork. *Journal of Food Protection*, 68, 1788e1798.

- 66 *In Food hygiene basic Texts* (4th ed.). Rome: WHO/FAO. Retrieved 10.12.2021 from www.fao.org/docrep/012/a1552e/a1552-00.pdf
- 67 Blagojevic, B., Antic, D., Ducic, M., Buncic, S. (2012). Visual cleanliness scores of cattle at slaughter and microbial loads on the hides and the carcasses. *Vet. Rec.*, 170, 563-570.
- 68 Madden, RH, Murray, KA, Gilmour, A. (2004). Determination of the principal points of product contamination during beef carcass dressing processes in Northern Ireland. *J. Food Prot.*, 67, 1494-1496.
- 69 Roberts, TA. (1980). Contamination of meat: the effects of slaughter practices on the bacteriology of the red meat carcass. *Journal of Royal Society of Health*, 100, 3-9.
- 70 Global GAP. 2020. Generic HACCP Food Safety Plan for crops-General guidance Version 5.4, Global G.A.P Foodplus, Köln.
- 71 Nørrung, B., Buncic, S. (2008). Microbial safety of meat in the European Union. *Meat Science*, 78(1-2), 14-24.
- 72 Nortjé, GL, Nel, L., Jordaan, E., Naudé, RT, Holzapfel, WH, Grimbeek, RJ. (1989a). A microbiological survey of fresh meat in the supermarket trade. Part 1: Carcasses and contact surfaces. *Meat Science*, 25(2), 81-97.
- 73 Igwaran, A., Okoh, AI. (2019). Human campylobacteriosis: A public health concern of global importance, *Heliyon*, 5:11, e02814. DOI: 10.1016/j.heliyon.2019.e02814.
- 74 Roberts, H., Jager, L., Blight, G. (2009). Waste-handling practices at red meat abattoirs in South Africa. *Waste Management & Research*, 27: 1, 25-30.
- 75 Adetunde, L., Glover, R., Oliver, A., Samuel, T. (2011). Source and distribution of microbial contamination on beef and Chevron in Navrongo, Kassena Nankana district of Upper East region in Ghana. *Journal of Animal Production Advances*, 1: 1, 21-28.
- 76 Haileselassie, M., Taddele, H., Adhana, K., Kalayou, S. (2013). Food safety knowledge and practices of abattoir and butchery shops and the microbial profile of meat in Mekelle city, Ethiopia. *Asian Pacific Journal of Tropical Biomedicine*, 3:5, 407-412.
- 77 Fasanmi, G., Olukole, S., Kehinde, O. (2010). Microbial studies of table scrapings from meat stalls in Ibadan metropolis, Nigeria: implications on meat hygiene. *African Journal of Biotechnology*, 9: 21, 3158-3162.
- 78 Codex Alimentarius International Food Standards. General Principles of Food Hygiene CXC: 1-1969. (2020). Codex Alimentarius, Roma.
- 79 Sayed, AA. (2019). Bacteriological and molecular studies on antibiotic resistant *Escherichia coli* isolated from meat and its products in Kaliobia, Egypt. *Benha Vet. Med.* 36(2), 335-344. DOI: 10.21608/BVMJ.2019.15531.1058.
- 80 Kuzuoka, K., Kawai, K., Yamauchi, S., Okada, A., Inoshima, Y. (2020). *Escherichia coli* O157 prevalence and enumeration of aerobic bacteria, Enterobacteriaceae, and *Escherichia coli* O157 at various steps in commercial beef processing plants, *Food Control*, 118, December 2020, 107353, DOI:10.1016/j.foodcont.2020.107353.
- 81 Prache, S., Adamiec, C., Astruc, T., Baéza-Campone, E., Bouillot, P., Clinquart, A., Feidt, C., Fourat, E., Gautron, J., Girard, A., Guillier, L., Kesse-Guyot, E., Lebreton, B., Lefèvre, F Le, Perchec, S., Martin, B., Mirade, P., Pierre, F., Raulet, M., Rémond, D., Sans, P., Souchon, I., Donnars, C. (2022) Review: Quality of animal-source foods. *Animal*, 1, 100376. DOI: 10.1016/j.animal.2021.100376.
- 82 Birhanu, W., Weldegebriel, S., Bassazin, G., Mitku, F., Birku, L., Tadesse, M. (2017). Assessment of microbiological quality and meat handling practices in butcher shops and abattoir found in Gondar town, Ethiopia. *International Journal of Microbiological Research*, 8:2, 59-68.
- 83 Bell, RG, Hathaway, SC. (1996). The hygienic efficiency of conventional and inverted lamb dressing systems. *Journal of Applied Bacteriology*, 81:3, 225-234.
- 84 Nørrung, B., Anderson, JK, Buncic, S. (2009). Main concerns of pathogenic microorganisms in meat. in *Safety of Meat and Processed Meat*, F. Toldra, Ed., Springer, New York, 3-29.
- 85 Song, X., Wang, H., Xu. X. (2021). Investigation of microbial contamination in a chicken slaughterhouse environment. *Journal of Food Science*, 86(8), 3598-3610. DOI: 10.1111/1750-3841.15842.
- 86 Fredriksson-Ahomaa, M., Gerhardt, M., Stolle, A. (2009). High bacterial contamination of pig tonsils at slaughter. *Meat Science*, 83, 334.

87 Jakubowska-Gawlik, K., Kolanowski, W., Murali, AP, Trafialek, J. (2022). A comparison of food safety conformity between cattle and pig slaughterhouses, *Food Control*, 109143.

89 Gill, CO, Deslandes, B., Rahn, K., Houde, A., Bryant, J. (1998). Evaluation of the hygienic performances of the processes for beef carcass dressing at 10 packing plants. *Journal of Applied Microbiology*, 84: 6, 1050-1058.

90 Ferreira Barros, M de A, Nero, LA, Monteiro AA, Beloti, V. (2007). Identification of main contamination points by hygiene indicator microorganisms in beef processing plants, *Ciênc. Tecnol. Aliment.*, 27:4, 856-862.

91 The two top zoonotic foodborne diseases in Europe in 2019 were caused by *Campylobacter*, with reported 220,682 cases (incidence of 59.7/100,000) and *Salmonella* with reported 87,923 cases (20.0/100,000). (2021). EFSA/ECDC.

92 Jenson, I., Sumner, J. (2012). Performance standards and meat safety - developments and direction. *Meat Science*, 92(3), 260-266.

93 Akaike, H., Nagai, M., Okatani, AT, Morita, Y. (2022). Food safety, livestock health, and productivity of a dairy farm following implementation of a certificated HACCP system. *Vet Med Sci.*, 84, 924-928. DOI: 10.1292/jvms.

94 Ministry of Agriculture Forestry and Fisheries. (04.01.2023). The improvement of the breeding hygiene management standard in the stage of production of the domestic animal, http://www.maff.go.jp/j/syouan/douei/katiku_yobo/k_haccp/index.html

95 WHO, Regional Office for Africa. (2004). Developing and maintaining food safety control systems for Africa current status and prospects for change, in Proceedings of the Second FAO/ WHO Global Forum of Food Safety Regulators, Bangkok, Thailand.

96 Birhanu, W., Weldegebriel, S., Bassazin, G., Mitku, F., Birku, L., Tadesse, M. (2017). Assessment of microbiological quality and meat handling practices in butcher shops and abattoir found in Gondar town, Ethiopia. *International Journal of Microbiological Research*, 8:2, 59-68.

97 Ortiz-Barrios, M., Miranda-De la Hoz, C., Lopez-Meza, P., Petrillo, A., De Felice, F. (2020). A case of food supply chain management with AHP, DEMATEL, and TOPSIS. *Journal of Multi-Criteria Decision Analysis*, 27, 104-128.

98 Nilda, TP, Rhamadani, A., Rhamadani, A., Wisnel, W. (2020). Designing food safety standards in beef jerky production process with the application of hazard analysis critical control point (HACCP), *Nutrition and Food Science*, 50, 333-347.

99 Hutchison, ML, Thomas, DJI, Small, AH, Buncic, S., Howell, M. (2007). Implementation of compulsory hazard analysis critical control point system and its effect on concentrations of carcass and environmental surface bacterial indicators in United Kingdom red meat slaughterhouses. *Journal of Food Protection*, 70(7), 1633-1639.

100 Nastasijevic, I., Mitrovic, R., Popovic, K., Tubic, M., Buncic, S. (2009). *The effects of a non-intervention HACCP implementation on process hygiene indicators on bovine and porcine carcasses*. Meso, XI (4), 232-239.

101 Tomasevic, I., Kuzmanović, J., Anđelković, A., Saračević, M., Stojanović, MM, Djekic, I. (2016). The effects of mandatory HACCP implementation on microbiological indicators of process hygiene in meat processing and retail establishments in Serbia. *Meat Science*, 114, 54-57. DOI: 10.1016/j.meatsci.2015.12.008].

References

1 Villarroel, A., Dargatz, DA, Lane, VM, McCluskey, BJ, Salman, MD. (2007). Suggested outline of potential critical control points for biosecurity and biocontainment on large dairy farms. *Food Control*, 203(6), 663-669. DOI: 10.2460/javma.230.6.808.

2 Sperber, WH. (2005). HACCP does not work from Farm to Table. *Food Control*, 16, 6, 511-514. DOI: 10.1016/j.foodcont.2003.10.013

3 Joint FAO/WHO Food Standards Program Codex Alimentarius Commission, Forty-sixth Session. FAO headquarters, Rome, Italy, 27 November to 2 December 2023.

4 Bae, YY, Choi, YM, Kim, MJ, Kim, KH, Kim, BC, Rhee, MS. (2011). Application of supercritical carbon dioxide for microorganism reductions in fresh pork. *J. Food Saf.*, 31, 511-517. DOI: 10.1111/j.1745-4565.2011.00328.x.

5 Kim, JH, Yim, DG. (2016). Assessment of the microbial level for livestock products in retail meat shops implementing HACCP system. *Korean J Food Sci An.*, 36, 594-600. DOI: 10.5851/kosfa.2016.36.5.594.

6 Agência Nacional de Vigilância Sanitária (2019). Instrução Normativa nº60 de 23 de dezembro de 2019, Estabelece as listas de padrões microbiológicos para alimentos. *Diário Oficial República Federativa do Brasil*, 249(1), 133.

7 Ribeiro Júnior, JC., Dias, BP, Nascimento, Nascimento, AL, Cabral Silva, JP, Cordeiro, RF, Oliveira Lobo, CM. (2024). Effects of washing sanitation standard operating procedures on the microbiological quality and safety of cattle carcasses. *Food Control*, 166. DOI: 10.1016/j.foodcont.2024.110745.

8 Clinquart, A., Ellies-Oury, P., Hocquette, F., Guillier, L., Santé-Lhoutellier, V., Prache, S. (2022). Review: On-farm and processing factors affecting bovine carcass and meat quality. *Animal*, 1, 100426. DOI: 10.1016/j.animal.2021.100426.

9 Kim, JH, Yim, DG. (2017). Microbial levels for food contact and environmental surfaces in meat processing plants and retail shops. *Food Sci Biotechnol.*, 26, 299-302.

10 Ahmed, AH, Al-Mahmood, OA. (2023) Food Safety Programs that should be Implemented in Slaughterhouses: Review. *Journal of Applied Veterinary Sciences*, 8(2), 80-88. DOI: 10.21608/JAVS.2023.185918.1208.

11 Wang, X., Bouzembrak, Y., Oude Lansink, AGJM, van der Fels-Klerx, HJ. (2021). Application of machine learning to the monitoring and prediction of food safety: A review. *Comprehensive Reviews in Food Science and Food Safety*, 21: 1, 416-434. DOI: 10.1111/1541-4337.

12 Abou El Nour-Mona M, Sakr-Ebtehag, AE. (2020). Isolation and Molecular. Identification of Some Food Borne Pathogens from Raw and Processed Meats: In vitro Synergistic Probability of Lactobacillus CFS and Antibiotic. *Egypt. J.Bot.*, 60(2), 405-421. DOI: 10.21608/EJBO.2019.19837.1395.

13 Kuzuoka, K., Kawai, K., Yamauchi, S., Okada, A., Inoshima, Y. (2020). Chilling control of beef and pork carcasses in a slaughterhouse based on causality analysis by graphical modelling. *Food Control*, 32, 106075. DOI: 10.1016/j.foodcont.2020.107353.

14 Prache, S., Schreurs, N., Guillier, L. (2022). Review: Factors affecting sheep carcass and meat quality attributes. *Animal*, 1:100330. DOI: 10.1016/j.animal.2021.100330.

15 Mansur, AR, Song, EJ, Cho, YS, Nam, YD, Choi, YS, Kim, DO. (2019). Comparative evaluation of spoilage-related bacterial diversity and metabolite profiles in chilled beef stored under air and vacuum packaging. *Food Microbiol*, 77, 166-72.

16 *European Surveillance of Veterinary Antimicrobial Consumption (ESVAC)*. (5.02.2023). <https://www.ema.europa.eu/en/veterinary-regulatory/overview/antimicrobial-resistance/european-surveillance-veterinary-antimicrobial-consumption-esvac>.

17 Abd El-Tawab, A., Maarouf, AA, El-Hofy-Fatma, I., Ahmed-Nesma, MG. (2020). Phenotypic characterization of some food poisoning bacteria isolated from meat and meat products in Kaliobia, Egypt. *Benha Vet. Med. J.*, 38(2), 146-151. DOI: 10.21608/bvmj.2020.25678.1186.

18 Røtterud, OJ, Gravning, GE, Hauge, SJ, Alvseike, O. (2020). Hygiene performancerating-an auditing scheme for evaluation of slaughter hygiene. *Methods X*, 20:7, 100829. DOI: 10.1016/j.mex.2020.100829.

19 Korany, SH, Tolba, KS, Sobhy, HM, Hekal, SA. (2021). Assuring the Safety of Local and Imported Beef Meat from Different Slaughterhouses in Egypt. *Advances in Animal and Veterinary Sciences*, 9(9). DOI: 10.17582/journal.aavs/2021/9.9.1472.1482.

20 Birhanu, W., Weldegebriel, S., Bassazin, G., Mitku, F., Birku, L., Tadesse, M. (2017). Assessment of microbiological quality and meat handling practices in butcher shops and abattoir found in Gondar town, Ethiopia. *International Journal of Microbiological Research*, 8:2, 59-68.

21 Food Control 140 (2022). 109143 8 FAO & WHO. (2020). COVID-19 and food safety: Guidance for food businesses. https://apps.who.int/iris/bitstream/handle/10665/331705/WHO-2019-nCoV-Food_Safety-2020.

22 Sagawa, R., Rodrigues, YM, Nascimento, CA, Ribeiro, J., Oliveira, MS, Muller, AC, Conti, ACM, Ribeiro Júnior, JC. (2021). Impact of the pre-slaughter period on the contamination of bovine leather and the operational sanitary procedure for skinning on the quality and microbiological safety of the carcass. *Semina: Ciênc. Agrár.*, 43, 4. DOI: 10.5433/1679-0359.2022v43n4p1835

23 Cavalcanti, AA, Limeira, CH, Siqueira, IN, de Lima, AC, de Medeiros, FJP, de Souza, JG, de Araújo Medeiros, NG, Abrahão Alves de, OF, de Melo, MA. (2022). The prevalence of *Listeria monocytogenes* in meat products in Brazil: A systematic literature review and meta-analysis. *Research in Veterinary Science*, 145, 169-176.

24 Perez-Rodriguez, F., Castro, R., Posada-Izquierdo, GD, Valero, A., Carrasco, E., Garcia-Gimeno, RM, Zurera, G. (2010). Evaluation of hygiene practices and microbiological quality of cooked meat products during slicing and handling at retail. *MeatSci.*, 86, 479-485. DOI: 10.1016/j.meatsci.2010.05.0387.

25 Alfian, G., Syafrudin M., Farooq, U., Ma'arif, MR, Syaekhoni, MA, Fitriyani, NL, Lee, J., Rhee, J. (2020). Improving efficiency of RFID-based traceability system for perishable food by utilizing IoT sensors and machine learning model. *Food Control*, 110, 107016. DOI: 10.1016/j.foodcont.2019.107016.

26 Cevallos-Almeida, M., Burgos-Mayorga, A., Gómez, CA, Lema-Hurtado, JL, Lema, L., Calvache, I., Jaramillo, C., Ruilova, IC, Martínez, EP, Estupiñán, P. (2021). Association between animal welfare indicators and microbiological quality of beef carcasses, including *Salmonella* spp., from a slaughterhouse in Ecuador. *Veterinary World*, 14(4), 918-925. DOI: 10.14202/vetworld.2021.918-925.

27 Oliveira, MS, Santos, IGC, Dias, BP, Nascimento, CA, Rodrigues, ÉM, Ribeiro, JC, Jr., Alfieri, AA, Alexandrino, B. (2021). Hygienic-health quality and microbiological hazard of clandestine Minas Frescal cheese commercialized in north Tocantins, Brazil. *Semina: Ciências Agrárias*, 42(2), 679-694. DOI: 10.5433/1679-0359.2021v42n2p679.

28 Birhanu, S., Menda, S. (2017). Hygienic Handling and Processing of Raw Beef Meat at Slaughter Houses and Meat Stalls in Gojjam Area. (*IJVHRS*), 5(8): 213-216. <https://doi.org/10.19070/2332-2748-1700042>.

29 Antic, D., Houf, K., Michalopoulou, E., Blagojevic, B. (2021). Beef abattoir interventions in a risk-based meat safety assurance system. *Meat Science*, 182, 108622. DOI: 10.1016/j.meatsci.2021.108622.

30 Nakamura, A., Takahashi, H., Kondo, A., Koike, F., Kuda, T., Kimura, B., Kobayashi, M. (2022). Distribution of psychrophilic microorganisms in a beef slaughterhouse in Japan after cleaning. *PLoS ONE*, 17(8), e0268411. DOI: 10.1371/journal.pone.0268411.

31 Antic, D., Houf, K., Michalopoulou, E., Blagojevic, B. (2021). Beef abattoir interventions in a risk-based meat safety assurance system. *Meat Science*, 182, 108622. DOI: 10.1016/j.meatsci.2021.108622.

32 Gutema, FD, Abdi, RD, Agga, GE, Firew, S., Rasschaert, G., Mattheus, W., Crombe, F., Duchateau, L., Gabriël, S., De Zutter, L. (2021). Assessment of beef carcass contamination with *Salmonella* and *E. coli* O 157 in slaughterhouses in Bishoftu, Ethiopia. *International Journal of Food Contamination*, 8(1), 1-9. DOI: 10.1186/s40550-021-00082-1.

33 De Filippis, F, Storia, AL, Villani, F, Ercolini, D. (2019). Strain-level diversity analysis of *Pseudomonas fragi* after in situ pangenome reconstruction shows distinctive spoilage associated metabolic traits clearly selected by different storage conditions. *Appl Environ Microbiol*, 85(1).

34 Carrasco-García, AA, Pardío-Sedas, VT, León-Banda, GG, Ahuja-Aguirre, C., Paredes-Ramos, P., Hernández-Cruz, BC, Murillo, VV. (2020). Effect of stress during slaughter on carcass characteristics and meat quality in tropical beef cattle. *Asian-Australasian Journal of Animal Sciences*, 33(10), 1656-1665. DOI: 10.5713/ajas.19.0804.

35 Nakamura, A., Takahashi, H., Koike, F., Kuda, T., Kobayashi, M. (2023). Transition of microbial contamination on the surface of carcass during the cattle slaughter process. *Food Microbiology*, 104245.

36 Calle, A., Carrascal, AK, Patiño, C., Carpio, C., Echeverry, A., Brashears, M. (2021). Seasonal effect on *Salmonella*, Shiga toxin-producing *E. coli* O157: H7 and non-O157 in the beef industry in Colombia, South America. *Heliyon*, 7(7), e07547. DOI: 10.1016/j.heliyon.2021.

37 Gill, CO. (2007). Microbiological conditions of meats from large game animals and birds. *Meat Science*, 77, 149-160.

38 Durmuşoğlu, H., İncili, GK, Demir, P, İlhak, Oİ. (2020). Effects of workers' hand washing and knife disinfection practices on microbiological quality of small animal carcasses in slaughterhouse environment. *Food Process Preserv.* 44(12), 1-7.

- 39 Kuzuoka, K., Kawai, K., Yamauchi, S., Okada, A., Inoshima, Y. (2018). Chilling control of beef and pork carcasses in a slaughterhouse based on causality analysis by graphical modelling. *Food Control*, 118, 107353.
- 40 Nakamura, A., Takahashi, H., Otomo, K., Mizuno, Y., Kuda, T., Kimura, B., Koike, F., Kobayashi, M. (2021). Dynamics of microbiota in Japanese Black beef stored for a long time under chilled conditions. *Food Microbiol*, 100, 103849.
- 41 Akaike, H., Nagai, M., Okatani, AT. Morita, Y. (2023). Food safety, livestock health, and productivity of a farm fattening Japanese black cattle following implementation of a certificated Hazard Analysis and Critical Control Point (HACCP) system. *J Vet Med Sci.*, 7:85(9), 985-989.
- 42 Alvseike, O., Røssvoll, E., Røtterud, OJ, Nesbakken, T., Skjerve, E., Prieto, M., Sandberg, M., Johannessen, G., Økland, M., Urdahl, AM, Hauge, SJ. (2019). Slaughter hygiene in European cattle and sheep abattoirs assessed by microbiological testing and Hygiene Performance Rating. *Food Control*, 101, 233-240.
- 43 Mallhi, IY, Sohaib, M., Khan, AU, Nawaz, M. (2019). Evaluating food safety knowledge, practices, and microbial profile of meat in abattoirs and butchery shops in Lahore, Pakistan. *Journal of Food Safety*, 39(5).
- 44 Odeyemi, OA, Alegbeleye, OO, Strateva, M, Stratev, D. (2020). Understanding spoilage microbial community and spoilage mechanisms in foods of animal origin. *Compr Rev Food Sci Food Saf.*, 19(2), 311-31.
- 45 Webb, NB, Marsden, JL. (1995). *Relationship of the HACCP system to Total Quality Management*. In: Person, AM, Dutson, TR. (Eds.), *Blackie Academic*, London, 10, 156-181.
- 46 Blagojević, B., Nesbakken, T., Alvseike, O., Vågsholm, I., Antic, D., Johlerf, S., Houf, K., Meemken, D., Nastasijevic, I., Vieira Pinto, M., Antunowic, B., Georgiev, M., Alban, L. (2021). Challenges and opportunities in the implementation of new meat inspection systems in Europe. *Trends in Food Science & Technology*, 116, 460-467.
- 47 Panea, B., Ripoll, G. (2020). Designing food safety standards in beef jerky production process with the application of hazard analysis critical control point (HACCP). *Nutrition & Food Science*, 50, 333-347.
- 48 Elder, RO, Keen, JE, Siragusa, GR, Barkocy-Gallagher, GA, Koohmaraie, M., Laegreid, WW. (2000). Correlation of Enterohemorrhagic Escherichia coli O157 prevalence in feces, hides, and carcasses of beef cattle during processing. *Proc. Natl. Acad. Sci.*, 97(7), 2999-3003.
- 49 Antic, D., Houf, K., Michalopoulou, E., Blagojevic, B. (2000). Beef abattoir interventions in a risk-based meat safety assurance system. *Meat Sci.*, 182, 108622. DOI: 10.1016/j.meatsci.2021.108622.
- 50 MIG, Microbiological Criteria Information for the UK Meat Industries, MIG, Moscow, Russia, 2006.
- 51 Ghafir, Y., China, B., Dierick, K., Zutter, LD, Daube, G. (2008). Hygiene indicator microorganisms for selected pathogens on beef, pork, and poultry meats in Belgium. *J. Food Prot.* 71(1), 35-45.
- 52 Gill, CO, Deslandes, B., Rahn, K., Houde, A., Bryant, J. (1998). Evaluation of the hygienic performances of the processes for beef carcass dressing at 10 packing plants. *J. Appl. Microbiol.* 84(6), 1050-1058.
- 53 Eisel, WG, Linton, RH, Muriana, PM. (1997). A survey of microbial levels for incoming raw beef, environmental sources, and ground beef in a red meat processing plant. *Food Microbiol.* 14:3, 273-282.
- 54 Wray, C., Davies, RH. (2000). *Salmonella in domestic animals*. CABI Publishing, Wallingford, UK. 184.
- 55 Lynn, TV, Hancock, DD, Besser, TE, Harrison, JH, Rice, DH, Stewart, NT, Rowan, LL. (1998). The occurrence and replication of Escherichia coli in cattle feeds. *J. Dairy Sci.*, 81(4), 1102-1108.
- 56 U.S. Department of Agriculture. (1997). USDA-APHIS: factors associated with Escherichia coli O157 in feces of feedlot cattle. Centers for Epidemiology and Animal Health, Fort Collins, Colo.
- 57 Armstrong, GL, Hollingsworth, J., Morris, JG Jr. (1996). Emerging foodborne pathogens: Escherichia coli O157:H7 as a model of entry of a new pathogen into the food supply of the developed world. *Epidemiol. Rev.*, 18(1), 29-51.

58 ANZFS (Australia New Zealand Food Standards Code) Microbiological Limits for Food - Schedule 27 - F2021C00605 (2021).

59 Laegreid, WW, Elder, RO, Keen, JE. (1999). Prevalence of *Escherichia coli* O157:H7 in range beef calves at weaning. *Epidemiol. Infect.*, 123, 291-298.

60 Cooper, K. (1999). The Plant Environment Counts: Protect your Product through Environmental, Sampling. *Meat and Poultry*, 5, 5.

61 Sofos, JN, Kochevar, SL, Bellinger, GR, Buege, DR, Hancock, DD, Ingham, SC, Morgan, JB, Reagan, JO, Smith, GC. (1999). Sources and extent of microbiological contamination of beef carcasses in seven United States slaughtering plants. *J Food Prot*, 62(2), 140-5.

62 Sudhakar, B, Paturkar, Waskar, VS, Zende, RJ. (2009). Bacteriological screening of environmental sources of contamination in an abattoir and the meat shops in Mumbai, India. *As. J. Food Ag-Ind.*, 2(3), 277-287.

63 Ali, N., Farooqui, A., Khan, A., Kazmi, S. (2010). Microbial contamination of raw meat and its environment in retail shops in Karachi, Pakistan. *Journal of Infection in Developing Countries*, 4: 6, 382-388.

64 Arquias, RK, Seixas, DB. (2021). Riscos do consumo de carne não inspecionada e as principais características sensoriais analisadas pelos consumidores-revisão de literatura. Universitário FG. <https://repositorio.animaeducacao.com.br/handle/ANIMA/13584>.

65 Miller, GY, Liu, X., McNamara, PE, Barber, DA. (2005). The influence of *Salmonella* in pigs preharvest and during pork processing on human health costs and risks from pork. *Journal of Food Protection*, 68, 1788e1798.

66 *In Food hygiene basic Texts* (4th ed.). Rome: WHO/FAO. Retrieved 10.12.2021 from www.fao.org/docrep/012/a1552e/a1552-00.pdf

67 Blagojevic, B., Antic, D., Ducic, M., Buncic, S. (2012). Visual cleanliness scores of cattle at slaughter and microbial loads on the hides and the carcasses. *Vet. Rec.*, 170, 563-570.

68 Madden, RH, Murray, KA, Gilmour, A. (2004). Determination of the principal points of product contamination during beef carcass dressing processes in Northern Ireland. *J. Food Prot.*, 67, 1494-1496.

69 Roberts, TA. (1980). Contamination of meat: the effects of slaughter practices on the bacteriology of the red meat carcass. *Journal of Royal Society of Health*, 100, 3-9.

70 Global GAP. 2020. Generic HACCP Food Safety Plan for crops-General guidance Version 5.4, Global G.A.P Foodplus, Köln.

71 Nørrung, B., Buncic, S. (2008). Microbial safety of meat in the European Union. *Meat Science*, 78(1-2), 14-24.

72 Nortjé, GL, Nel, L., Jordaan, E., Naudé, RT, Holzapfel, WH, Grimbeek, RJ. (1989a). A microbiological survey of fresh meat in the supermarket trade. Part 1: Carcasses and contact surfaces. *Meat Science*, 25(2), 81-97.

73 Igwaran, A., Okoh, AI. (2019). Human campylobacteriosis: A public health concern of global importance, *Heliyon*, 5:11, e02814. DOI: 10.1016/j.heliyon.2019.e02814.

74 Roberts, H., Jager, L., Blight, G. (2009). Waste-handling practices at red meat abattoirs in South Africa. *Waste Management & Research*, 27: 1, 25-30.

75 Adetunde, L., Glover, R., Oliver, A., Samuel, T. (2011). Source and distribution of microbial contamination on beef and Chevron in Navrongo, Kassena Nankana district of Upper East region in Ghana. *Journal of Animal Production Advances*, 1: 1, 21-28.

76 Haileselassie, M., Taddele, H., Adhana, K., Kalayou, S. (2013). Food safety knowledge and practices of abattoir and butchery shops and the microbial profile of meat in Mekelle city, Ethiopia. *Asian Pacific Journal of Tropical Biomedicine*, 3:5, 407-412.

77 Fasanmi, G., Olukole, S., Kehinde, O. (2010). Microbial studies of table scrapings from meat stalls in Ibadan metropolis, Nigeria: implications on meat hygiene. *African Journal of Biotechnology*, 9: 21, 3158-3162.

78 Codex Alimentarius International Food Standards. General Principles of Food Hygiene CXC: 1-1969. (2020). Codex Alimentarius, Roma.

- 79 Sayed, AA. (2019). Bacteriological and molecular studies on antibiotic resistant *Escherichia coli* isolated from meat and its products in Kaliobia, Egypt. *Benha Vet. Med.* 36(2), 335-344. DOI: 10.21608/BVMJ.2019.15531.1058.
- 80 Kuzuoka, K., Kawai, K., Yamauchi, S., Okada, A., Inoshima, Y. (2020). *Escherichia coli* O157 prevalence and enumeration of aerobic bacteria, Enterobacteriaceae, and *Escherichia coli* O157 at various steps in commercial beef processing plants, *Food Control*, 118, December 2020, 107353, DOI:10.1016/j.foodcont.2020.107353.
- 81 Prache, S., Adamiec, C., Astruc, T., Baéza-Campone, E., Bouillot, P., Clinquart, A., Feidt, C., Fourat, E., Gautron, J., Girard, A., Guillier, L., Kesse-Guyot, E., Lebret, B., Lefèvre, F Le, Percec, S., Martin, B., Mirade, P., Pierre, F., Raulet, M., Rémond, D., Sans, P., Souchon, I., Donnars, C. (2022) Review: Quality of animal-source foods. *Animal*, 1, 100376. DOI: 10.1016/j.animal.2021.100376.
- 82 Birhanu, W., Weldegebriel, S., Bassazin, G., Mitku, F., Birku, L., Tadesse, M. (2017). Assessment of microbiological quality and meat handling practices in butcher shops and abattoir found in Gondar town, Ethiopia. *International Journal of Microbiological Research*, 8:2, 59-68.
- 83 Bell, RG, Hathaway, SC. (1996). The hygienic efficiency of conventional and inverted lamb dressing systems. *Journal of Applied Bacteriology*, 81:3, 225-234.
- 84 Norrung, B., Anderson, JK, Buncic, S. (2009). *Main concerns of pathogenic microorganisms in meat.* in Safety of Meat and Processed Meat, F. Toldra, Ed., Springer, New York, 3-29.
- 85 Song, X., Wang, H., Xu. X. (2021). Investigation of microbial contamination in a chicken slaughterhouse environment. *Journal of Food Science*, 86(8), 3598-3610. DOI: 10.1111/1750-3841.15842.
- 86 Fredriksson-Ahomaa, M., Gerhardt, M., Stolle, A. (2009). High bacterial contamination of pig tonsils at slaughter. *Meat Science*, 83, 334.
- 87 Jakubowska-Gawlik, K., Kolanowski, W., Murali, AP, Trafialek. J. (2022). A comparison of food safety conformity between cattle and pig slaughterhouses, *Food Control*, 109143.
- 89 Gill, CO, Deslandes, B., Rahn, K., Houde, A., Bryant, J. (1998). Evaluation of the hygienic performances of the processes for beef carcass dressing at 10 packing plants. *Journal of Applied Microbiology*, 84: 6, 1050-1058.
- 90 Ferreira Barros, M de A, Nero, LA, Monteiro AA, Beloti, V. (2007). Identification of main contamination points by hygiene indicator microorganisms in beef processing plants, *Ciênc. Tecnol. Aliment.*, 27:4, 856-862.
- 91 The two top zoonotic foodborne diseases in Europe in 2019 were caused by *Campylobacter*, with reported 220,682 cases (incidence of 59.7/100,000) and *Salmonella* with reported 87,923 cases (20.0/100,000). (2021). EFSA/ECDC.
- 92 Jenson, I., Sumner, J. (2012). Performance standards and meat safety - developments and direction. *Meat Science*, 92(3), 260-266.
- 93 Akaike, H., Nagai, M., Okatani, AT, Morita, Y. (2022). Food safety, livestock health, and productivity of a dairy farm following implementation of a certificated HACCP system. *Vet Med Sci.*, 84, 924-928. DOI: 10.1292/jvms.
- 94 Ministry of Agriculture Forestry and Fisheries. (04.01.2023). The improvement of the breeding hygiene management standard in the stage of production of the domestic animal, http://www.maff.go.jp/j/syouan/douei/katiku_yobo/k_haccp/index.html
- 95 WHO, Regional Office for Africa. (2004). Developing and maintaining food safety control systems for Africa current status and prospects for change, in Proceedings of the Second FAO/ WHO Global Forum of Food Safety Regulators, Bangkok, Thailand.
- 96 Birhanu, W., Weldegebriel, S., Bassazin, G., Mitku, F., Birku, L., Tadesse, M. (2017). Assessment of microbiological quality and meat handling practices in butcher shops and abattoir found in Gondar town, Ethiopia. *International Journal of Microbiological Research*, 8:2, 59-68.
- 97 Ortiz-Barríos, M., Miranda-De la Hoz, C., Lopez-Meza, P., Petrillo, A., De Felice, F. (2020). A case of food supply chain management with AHP, DEMATEL, and TOPSIS. *Journal of Multi-Criteria Decision Analysis*, 27, 104-128.
- 98 Nilda, TP, Rhamadani, A., Rhamadani, A., Wisnel, W. (2020). Designing food safety standards in beef jerky production process with the application of hazard analysis critical control point (HACCP), *Nutrition and Food Science*, 50, 333-347.

99 Hutchison, ML, Thomas, DJI, Small, AH, Buncic, S., Howell, M. (2007). Implementation of compulsory hazard analysis critical control point system and its effect on concentrations of carcass and environmental surface bacterial indicators in United Kingdom red meat slaughterhouses. *Journal of Food Protection*, 70(7), 1633-1639.

100 Nastasijevic, I., Mitrovic, R., Popovic, K., Tubic, M., Buncic, S. (2009). *The effects of a non-intervention HACCP implementation on process hygiene indicators on bovine and porcine carcasses*. Meso, XI (4), 232-239.







101 Tomasevic, I., Kuzmanović, J., Anđelković, A., Saračević, M., Stojanović, MM, Djekic, I. (2016). The effects of mandatory HACCP implementation on microbiological indicators of process hygiene in meat processing and retail establishments in Serbia. *Meat Science*, 114, 54-57. DOI: 10.1016/j.meatsci.2015.12.008].

Herald of Science of S.Seifullin Kazakh Agrotechnical Research University: Veterinary Sciences.
– Astana: S. Seifullin Kazakh Agrotechnical Research University, 2024. – № 4 (008). – P. 34-42.
- ISSN 2958-5430, ISSN 2958-5449

doi.org/ 10.51452/kazatuvc.2024.4(008).1798
UDC579.672

Research article

Analysis of microbial contamination in different types of meat in the Kostanay region

Raushan M. Rychshanova¹ , Anara M. Mendybayeva² , Yuliya Y. Aleshina¹ ,
Madina S. Ernazarova¹ , Zulkiya B. Abilova¹ , Albina R. Gabitova¹ 

¹Akhmet Baitursynuly Kostanay Regional University, Kostanay, Kazakhstan

²"Antigen" Scientific and Production Enterprise LLP, Almaty, Kazakhstan

Corresponding author: Albina R. Gabitova: bibishka03@mail.ru

Co-authors: (1: RR) raushan5888@mail.ru; (2: AM) jks1992@mail.ru;

(3: YuA) juliya.240895@gmail.com; (4: ME) madina.ernazarova@mail.ru;

(5: ZA) dgip2005@mail.ru

Received: 24-10-2024 **Accepted:** 17-12-2024 **Published:** 30-12-2024

Abstract

Background and Aim. Meat is a favourable environment for the growth and viability of pathogenic microorganisms. Bacteria in meat cause spoilage and increase the risk of foodborne toxic infections of various origins during consumption. The importance of monitoring and analysing microbial contamination is increasing due to increased requirements for food safety and public health protection. The aim of the study was to analyse the overall level of microbial contamination of different types of meat sampled from retail outlets in the Kostanay region.

Materials and Methods. A total of 30 meat samples, including pork, beef, and horse meat, were collected for analysis. To assess microbial contamination of the meat, the number of mesophilic aerobic and facultatively anaerobic microorganisms (QMAFAnM) and their species composition were determined.

Results. It was found that 96% of the meat sold at the retail level had elevated QMAFAnM, with coliforms isolated in 30% of the samples. *Salmonella spp.* and *Listeria monocytogenes* were not detected in the meat tested. Pork meat had the highest microbial contamination compared to beef and horse meat samples.

Conclusion. Exceeding the permitted levels of QMAFAnM and the presence of coliform bacteria indicate potential health risks for consumers and more effective measures are required to ensure food safety.

Keywords: coliforms; food safety; meat; microbial contamination; QMAFAnM.

Introduction

Food safety and consumer protection are fundamental priorities in government policies worldwide. Ensuring the safety of raw materials and food products is essential for maintaining public health. Food quality directly impacts population health, helping to reduce the risk of infectious and chronic diseases. This underscores the need for strict control over the production, storage, and distribution of food products.

Meat is an important component of a healthy diet, rich in high-quality protein essential for the normal functioning of all body systems [1]. The proteins in meat provide all the essential amino acids necessary for the growth and repair of tissues. Additionally, meat contains B-group vitamins, which play crucial roles in energy metabolism and various metabolic processes, helping maintain a healthy nervous system and cognitive abilities [2]. Meat is also abundant in micronutrients like magnesium, selenium, and zinc that support the immune system and are involved in producing hormones and enzymes [3].

Meat also supplies bioavailable forms of iron and phosphorus, which are essential for oxygen transport and bone mineralization, respectively [4]. The unique nutrient composition of meat makes it a key dietary component for sustaining cellular functions and biochemical processes critical to physiological health.

However, it should be noted that meat and meat products can potentially source several infectious diseases. In particular, meat derived from animals afflicted with illness has the capacity to result in human infection with zoonotic diseases [5, 6].

The contamination of meat with microbial organisms represents a significant challenge to consumer health and the quality of food products in the livestock and food industries. Pathogenic bacteria, including *Escherichia coli*, *Listeria monocytogenes* and *Staphylococcus aureus*, may be present in poultry, pork, mutton and beef. Shiga-toxin-producing *E. coli* is frequently associated with beef and is responsible for a range of severe toxic infections, including haemorrhagic colitis and haemolytic uremic syndrome [7]. *L. monocytogenes* is a facultative anaerobic Gram-positive bacterium that is found in fresh meat and meat products. However, human listeriosis is mainly associated with ready-to-eat meat products with a long shelf life in the refrigerator [8]. A prevalence of 35% was observed for *S. aureus* in retail meat samples collected in 39 cities in China between 2011 and 2016 [7].

Furthermore, the contravention of storage conditions gives rise to the deterioration of foodstuffs and an augmented probability of contracting food poisoning upon consumption [7, 9]. Statistical evidence indicates that meat spoilage is responsible for approximately 400,000 deaths and over 600 million disease cases annually. The World Health Organization (WHO) has reported that a significant proportion of the global population has experienced food-related illness at some point in their lives, with over 30 diseases linked to poor food quality [10]. The relevance of this issue is also confirmed by data from Kazakhstan. In 2020, 37 cases of acute intestinal infections per 100,000 people were recorded, while in 2023, this figure increased to 55.37 cases per 100,000 people. The rise in morbidity highlights the need to tighten control over the quality of food products and improve sanitary and hygienic conditions at all stages of meat production and distribution [11].

The aim of our research is to analyse the overall level of microbial contamination of meat of different types of animals sampled from retail outlets in Kostanay region.

Materials and Methods

The study was conducted in the microbiological analysis laboratory of the Scientific Research Institute of Applied Biotechnology at A. Baitursynuly Kostanay Regional University. A total of 30 samples of chilled meat, including pork (n=10), beef (n=10), and horse meat (n=10), were collected from five different retail outlets in accordance with the GOST 31904-2012 standard “Food products. Methods of sampling for microbiological tests” [12].

Samples were delivered to the laboratory in sterile containers, with temperature control maintained to preserve sample integrity for microbiological analysis. Upon arrival, the samples were processed immediately under aseptic conditions to prevent any extraneous contamination.

The microbiological methods employed in the study were conducted in accordance with the standards outlined in GOST [13-18].

For the identification of coliform bacteria, samples were placed in meat-peptone broth (FBIS SRCAMB, Obolensk, Russia), followed by subculturing onto Endo agar and TBX agar (CHROMagar, Paris, France) and incubated at 37 °C for 24 hours. Colonies with characteristic morphology were subjected to further biochemical analysis.

For *S. aureus*, yolk-salt agar was used, with incubation at 37 °C for 24 hours, and then at room temperature for an additional 24 hours, with confirmatory tests based on coagulase activity.

Detection of *Salmonella* spp. included pre-enrichment in buffered peptone water (FBIS SRCAMB, Obolensk, Russia), selective enrichment in Rappaport-Vassiliadis broth (FBIS SRCAMB, Obolensk, Russia), and subsequent inoculation onto Salmonella agar (CHROMagar, Paris, France).

To identify *L. monocytogenes*, samples were placed in Fraser broth (FBIS SRCAMB, Obolensk, Russia), enriched, and incubated for two days, with the first day at 30 °C and the second at 37 °C, followed by subculturing onto Listeria chromogenic agar (CHROMagar, Paris, France) and Oxford agar (Condalab, Madrid, Spain).

To determine the total microbial number (QMAFAnM), which is the quantity of mesophilic aerobic and facultative-anaerobic microorganisms, we conducted studies using the classical deep culture method in accordance with the standards outlined in GOST 26670-91 [18].

The presence of total microbial contamination was assessed by inoculating serial dilutions of the samples onto QMAFAnM media (FBIS SRCAMB, Obolensk, Russia) and incubating at 37° C for 24 hours, followed by an additional 24 hours at room temperature.

The arithmetic mean of the number of colonies from all cultures was calculated using the counting process results. The number of microorganisms in 1.0 g (“X”) of the product was calculated using the following formula:

$$X = n \times 10^m,$$

this formula defines the variables as follows: “n” represents the number of colonies counted on the dish, while “m” denotes the number of decimal dilutions.

The calculation result was expressed as a number between 1.0 and 9.9×10^m . The obtained number of microorganisms was then subjected to a verification process in accordance with the relevant requirements outlined in “Customs Union Technical Regulations on the safety of food products 021/2011” [19].

Results

Among the microbiological indicators, we conducted studies to determine the presence of coliform bacteria. Characteristic growth of *E. coli* was observed on solid differential Endo medium, forming colonies with a metallic sheen, and on CHROMagar TBX chromogenic medium, where colonies appeared blue. *Proteus spp.* colonies on Endo agar were transparent with swarming growth and a distinctive odor. Growth of *E. coli* (7 strains) and *Proteus spp.* (2 strains) was detected (Figure 1).





Figure 1 – Microbiological investigation of coliform bacteria

Research was also carried out to detect *S. aureus*. Staphylococci demonstrated characteristic growth on yolk-salt agar, forming colonies with “a rainbow halo” (indicating lecithinase production). A total of 4 strains coagulase-positive *S. aureus* were detected (Figure 2).



Growth of *S. aureus* in sample No. 26 on yolk-salt agar

Figure 2 – Microbiological investigation of *S. aureus*

The results of these studies demonstrated that *L. monocytogenes* and *Salmonella spp.* bacteria were not detected in any of the tested samples.

The number of colonies grown was enumerated in the cultures. The quantity of microbial colonies was determined by counting in cultures of dilutions at 10^2 , 10^3 , 10^4 , respectively (Figure 3).




Figure 3 – Microbiological investigation QMAFAnM (dilutions: 10^2 , 10^3 , 10^4)

The findings of the study indicated that all 10 pork samples analysed exceeded the permissible level of QMAFAnM. Coliforms were identified in four samples (No. 2, No.8 - *Proteus spp.*, No.5, No.7 - *E. coli*). Two strains of *S. aureus* were isolated from samples No. 2 and No. 8. However, no evidence of *Salmonella spp.* or *Listeria spp.* bacteria was observed (Table 1).

Table 1 – Results of microbiological examination of pork

Sample №	QMAFAnM, CFU/g	Coliforms	<i>Salmonella spp.</i>	<i>L. monocytogenes</i>	<i>S. aureus</i>
1	11×10^4	–	–	–	–
2	15×10^4	+	–	–	+
3	5.4×10^4	–	–	–	–
4	7.9×10^4	–	–	–	–
5	8.2×10^4	+	–	–	–
6	8.5×10^4	–	–	–	–
7	35×10^4	+	–	–	–
8	27×10^4	+	–	–	+
9	10×10^4	–	–	–	–
10	9.7×10^4	–	–	–	–
Norma	1×10^4 CFU/g, max.	Not allowed in 0.001 g of product	Not allowed in 25 g of product	Not allowed in 25 g of product	Not allowed in 1 g of product

The analysis results of beef samples indicate that all tested meat samples exceed the permissible level of QMAFAnM. The isolation of coliform bacteria was observed in three samples (No.12, No.16, No.20 - *E. coli*). The tested samples did not show the presence of *S. aureus*, *Salmonella*, or *Listeria species* (Table 2).

Table 2 – Results of microbiological examination of beef

Sample №	QMAFAnM, CFU/g	Coliforms	<i>Salmonella spp.</i>	<i>L. monocytogenes</i>	<i>S. aureus</i>
11	5.9×10^4	–	–	–	–
12	15×10^4	+	–	–	–
13	24×10^4	–	–	–	–
14	11×10^4	–	–	–	–
15	22×10^4	–	–	–	–
16	3.4×10^4	+	–	–	–
17	1.2×10^4	–	–	–	–
18	9.7×10^4	–	–	–	–
19	6.8×10^4	–	–	–	–
20	5.6×10^4	+	–	–	–
Norma	1×10^4 CFU/g, max.	Not allowed in 0.001 g of product	Not allowed in 25 g of product	Not allowed in 25 g of product	Not allowed in 1 g of product

The data obtained indicated that nine samples of horse meat under investigation had exceeded the maximum permitted level of contaminants as defined by the QMAFAnM. In two samples, a positive reaction for the presence of coliform was identified (samples No.21 and No.29, which tested positive for *E. coli*). Two strains of *S. aureus* were isolated from samples No. 26 and No.28. No evidence of *Salmonella* or *Listeria bacteria* was found in any of the tested samples (Table 3).

Table 3 – Results of microbiological examination of horse meat

Sample №	QMAFAnM, CFU/g	Coliforms	<i>Salmonella spp.</i>	<i>L. monocytogenes</i>	<i>S. aureus</i>
21	12×10^5	+	–	–	–
22	5.7×10^5	–	–	–	–
23	5.2×10^5	–	–	–	–
24	5.0×10^5	–	–	–	–
25	7.3×10^5	–	–	–	–
26	11.1×10^5	+	–	–	+
27	6.3×10^5	–	–	–	–
28	10.1×10^5	–	–	–	+
29	6.6×10^5	+	–	–	–
30	7.9×10^5	–	–	–	–
Norma	1×10^4 CFU/g, max.	Not allowed in 0.001 g of product	Not allowed in 25 g of product	Not allowed in 25 g of product	Not allowed in 1 g of product

The results of the studies demonstrate notable discrepancies in QMAFAnM levels across diverse meat categories. The results indicated that the mean QMAFAnM value in pork was 13.77×10^4 CFU/g, the highest among all the meat types under investigation. This suggests that pork is more susceptible to microbial contamination than other meats. Beef demonstrated a mean QMAFAnM value of 10.46×10^4 CFU/g, which is considerably lower than pork but higher than horse meat. Horse meat exhibited the lowest level of QMAFAnM, at 7.72×10^4 CFU/g, and is the least susceptible to microbial contamination among the tested meats (Figure 3).




Figure 3 – Indicators of the level of microbial contamination in the studied meat samples

Discussion and Conclusion

The study's results revealed notable differences in microbial contamination levels across different meat types. In all pork samples analyzed, the total microbial count (QMAFAnM) exceeded permissible limits, consistent with findings from other studies on microbial contamination in pork [20, 21]. Additionally, coliform bacteria and *S. aureus* were detected in pork samples, indicating potential health risks for consumers and possible lapses in hygiene protocols in the handling and processing of pork.

A similar contamination pattern was observed in the beef samples. All beef samples analyzed also had QMAFAnM levels above the acceptable standards, with coliforms present in three samples. This finding aligns with previous research [22] and may suggest deficiencies in the sanitary and hygienic processing of beef, potentially at various stages from production to storage. These issues highlight the need for more rigorous quality control measures for beef to prevent microbial contamination that could pose health risks.

The analysis of horsemeat samples showed that nine out of ten samples had QMAFAnM concentrations above permissible limits, though contamination levels were generally lower than those observed in pork and beef. Coliform bacteria and *S. aureus* were detected in two horsemeat samples, suggesting a degree of bacterial contamination but at a lower frequency than in the other meat types. This may reflect variations in the conditions under which horsemeat is handled and stored, yet underscores the importance of maintaining strict hygiene practices across all meat types.

Importantly, no samples from any of the meat types tested positive for the presence of pathogenic bacteria such as *Salmonella spp.* or *L. monocytogenes* indicating that while general bacterial contamination was an issue, there was no evidence of these specific pathogens in the samples.

To enhance the safety and quality of meat, adherence to microbiological control measures as set forth by the technical regulations of the Customs Union (021/2011,034/2013) [19, 23] is crucial. These regulations govern the production, storage, transportation, sale, and disposal of food products, ensuring safety standards are upheld at every stage in the supply chain.

The study results indicate the necessity for enhanced monitoring and regulation of the sanitary and hygienic conditions associated with the production, processing, and storage of meat of diverse types. The presence of elevated levels of QMAFAnM and the occurrence of coliform bacteria in meat products suggest potential risks to consumer health, underscoring the importance of implementing more rigorous measures to ensure food safety. Regular monitoring of microbial contamination and staff training in meat handling and storage practices are crucial strategies for reducing the risk of contamination.

Authors' Contribution

RR and AM: conceptualized and designed the study, conducted a comprehensive literature search, and analyzed the collected data. YuA, ME, and AG: carried out the research implementation and

analyzed the results. RR, ZA, and AG: performed the final editing and proofreading of the manuscript. All authors have read, reviewed, and approved the final version of the manuscript.

Acknowledgments

The study was conducted in accordance with the scientific and technical programme IRN BR22885795 "Improving Food Safety", which was funded by the Ministry of Agriculture of the Republic of Kazakhstan for the period 2024-2026.

References

- 1 Geiker, NRW, Bertram, HC, Mejborn, H., Dragsted, LO, Kristensen, L., Carrascal, JR, Bügel, S., Astrup, A. (2021). Meat and Human Health-Current Knowledge and Research Gaps. *Foods*, 10(7), 1556. DOI:10.3390/foods10071556.
- 2 Smith, NW, Fletcher, AJ, Hill, JP, McNabb, WC. (2022). Modeling the Contribution of Meat to Global Nutrient Availability. *Frontiers in nutrition*, 9, 766796. DOI:10.3389/fnut.2022.766796.
- 3 Narayanam, H., Chinni, SV, Samuggam, S. (2021). The Impact of Micronutrients-Calcium, Vitamin D, Selenium, Zinc in Cardiovascular Health: A Mini Review. *Frontiers in physiology*, 12, 742425. DOI:10.3389/fphys.2021.742425.
- 4 Falowo, A. (2021). A Comprehensive Review of Nutritional Benefits of Minerals in Meat and Meat Products. *Science Letters*, 9(2), 55-64. DOI:10.47262/SL/9.2.132021010.
- 5 Espinosa, R, Tago, D, Treich, N. (2020). Infectious Diseases and Meat Production. *Environ Resource Econ*, 76, 1019-1044. DOI:10.1007/s10640-020-00484-3.
- 6 Augustin, JC, Kooh, P, Bayeux, T, Guillier, L, Meyer, T, Jourdan-Da Silva, N, Villena, I, Sanaa, M., Cerf, O., on Behalf of the Anses Working Group on Consumer Information on Foodborne Biological Risks. (2020). Contribution of Foods and Poor Food-Handling Practices to the Burden of Foodborne Infectious Diseases in France. *Foods*, 9(11), 1644. DOI:10.3390/foods9111644.
- 7 Li, H., Sun, X., Liao, X., Gänzle, M. (2020). Control of pathogenic and spoilage bacteria in meat and meat products by high pressure: Challenges and future perspectives. *Comprehensive reviews in food science and food safety*, 19(6), 3476–3500. DOI:10.1111/1541-4337.12617.
- 8 Shao, L., Chen, S., Wang, H., Zhang, J., Xu, X., Wang, H. (2021). Advances in understanding the predominance, phenotypes, and mechanisms of bacteria related to meat spoilage. *Trends in Food Science & Technology*, 118, 822–832. DOI: 10.1016/j.tifs.2021.11.007.
- 9 Ashiq, J., Saeed, U., Li, Z., Nawaz, MH. (2024). Advances in meat spoilage detection: A review of methods involving 2D-based nanomaterials for detection of spoiled meat. *Journal of Food Composition and Analysis*, 132, 106295. DOI: 10.1016/j.jfca.2024.106295.
- 10 Matle, I., Mbatha, KR, Madoroba, E. (2020). A review of *Listeria monocytogenes* from meat and meat products: Epidemiology, virulence factors, antimicrobial resistance and diagnosis. *Onderstepoort Journal of Veterinary Research*, 87(1). DOI:10.4102/ojvr.v87i1.1869.
- 11 Агентство по стратегическому планированию и реформам Республики Казахстан Бюро национальной статистики. (2023). Статистика здравоохранения и социального обеспечения.
- 12 Межгосударственный совет по стандартизации, метрологии и сертификации. (2014). Межгосударственный стандарт: Продукты пищевые. Методы отбора проб для микробиологических испытаний (ГОСТ 31904-2012). Москва: Стандартинформ.
- 13 Межгосударственный совет по стандартизации, метрологии и сертификации. (2010). ГОСТ 26669-85. Межгосударственный стандарт: Продукты пищевые и вкусовые. Подготовка проб для микробиологического анализа (ГОСТ 26669-85). Москва: Стандартинформ.
- 14 Межгосударственный совет по стандартизации, метрологии и сертификации. (2013). Межгосударственный стандарт: Продукты пищевые. Методы выявления и определения количества бактерий группы кишечных палочек (колиформных бактерий) (ГОСТ 31747-2012). Москва: Стандартинформ.
- 15 Межгосударственный совет по стандартизации, метрологии и сертификации. (2014). Межгосударственный стандарт: Продукты пищевые. Метод выявления бактерий рода *Salmonella* (ГОСТ 31659-2012). Москва: Стандартинформ.

16 Межгосударственный совет по стандартизации, метрологии и сертификации. (2022). Межгосударственный стандарт: Продукты пищевые. Методы выявления бактерий *Listeria monocytogenes* и других видов *Listeria (Listeria spp.)* (ГОСТ 32031-2022). Москва: Стандартинформ.

17 Межгосударственный совет по стандартизации, метрологии и сертификации. (2013). Межгосударственный стандарт: Продукты пищевые. Методы выявления и определения количества коагулазоположительных стафилококков и *Staphylococcus aureus* (ГОСТ 31746-2012). Москва: Стандартинформ.

18 Межгосударственный совет по стандартизации, метрологии и сертификации. (2008). Межгосударственный стандарт: Продукты пищевые. Методы культивирования микроорганизмов (ГОСТ 26670-91). Москва: Стандартинформ.

19 Евразийская экономическая комиссия. (2011). Технический регламент Таможенного союза ТР ТС 021/2011. О безопасности пищевой продукции.

20 Ngo, ННТ, Nguyen-Thanh, L., Pham-Duc, P., Dang-Xuan, S., Le-Thi, H., Denis-Robichaud, J., Nguyen-Viet, H., Le, TT, Grace, D., Unger, F. (2021). Microbial contamination and associated risk factors in retailed pork from key value chains in Northern Vietnam. *International Journal of Food Microbiology*, 346, 109163. DOI: 10.1016/j.ijfoodmicro.2021.109163.

21 Veselá, H., Dorotíková, K., Dušková, M., Furmančíková, P., Šedo, O., Kameník, J. (2022). The pork meat or the environment of the production facility? The effect of individual technological steps on the bacterial contamination in cooked hams. *Microorganisms*, 10(6), 1106. DOI: 10.3390/microorganisms10061106.

22 Lenchenko, E., Stepanov, D. (2020). Quantitative account and differential properties of pathogenic bacteria allocated from food raw materials. *Problems of Veterinary Sanitation Hygiene and Ecology*, 1(2), 228–235 DOI: 10.36871/vet.san.hyg.ecol.202002017.

23 Евразийская экономическая комиссия. (2013). Технический регламент Таможенного союза ТР ТС 034/2013. О безопасности мяса и мясной продукции.

References

1 Geiker, NRW, Bertram, HC, Mejbörn, H., Dragsted, LO, Kristensen, L., Carrascal, JR, Bügel, S., Astrup, A. (2021). Meat and Human Health-Current Knowledge and Research Gaps. *Foods*, 10(7), 1556. DOI: 10.3390/foods10071556.

2 Smith, NW, Fletcher, AJ, Hill, JP, McNabb, WC. (2022). Modeling the Contribution of Meat to Global Nutrient Availability. *Frontiers in nutrition*, 9, 766796. DOI:10.3389/fnut.2022.766796.

3 Narayanam, H., Chinni, SV, Samuggam, S. (2021). The Impact of Micronutrients-Calcium, Vitamin D, Selenium, Zinc in Cardiovascular Health: A Mini Review. *Frontiers in physiology*, 12, 742425. DOI:10.3389/fphys.2021.742425.

4 Falowo, A. (2021). A Comprehensive Review of Nutritional Benefits of Minerals in Meat and Meat Products. *Science Letters*, 9(2), 55-64. DOI:10.47262/SL/9.2.132021010.

5 Espinosa, R, Tago, D, Treich, N. (2020). Infectious Diseases and Meat Production. *Environ Resource Econ*, 76, 1019-1044. DOI:10.1007/s10640-020-00484-3.

6 Augustin, JC, Kooh, P, Bayeux, T, Guillier, L, Meyer, T, Jourdan-Da Silva, N, Villena, I., Sanaa, M., Cerf, O., on Behalf of the Anses Working Group on Consumer Information on Foodborne Biological Risks. (2020). Contribution of Foods and Poor Food-Handling Practices to the Burden of Foodborne Infectious Diseases in France. *Foods*, 9(11), 1644. DOI:10.3390/foods9111644.

7 Li, H., Sun, X., Liao, X., Gänzle, M. (2020). Control of pathogenic and spoilage bacteria in meat and meat products by high pressure: Challenges and future perspectives. *Comprehensive reviews in food science and food safety*, 19(6), 3476–3500. DOI:10.1111/1541-4337.12617.

8 Shao, L., Chen, S., Wang, H., Zhang, J., Xu, X., Wang, H. (2021). Advances in understanding the predominance, phenotypes, and mechanisms of bacteria related to meat spoilage. *Trends in Food Science & Technology*, 118, 822–832. DOI: 10.1016/j.tifs.2021.11.007.

9 Ashiq, J., Saeed, U., Li, Z., Nawaz, MH. (2024). Advances in meat spoilage detection: A review of methods involving 2D-based nanomaterials for detection of spoiled meat. *Journal of Food Composition and Analysis*, 132, 106295. DOI: 10.1016/j.jfca.2024.106295.

10 Matle, I., Mbatha, KR, Madoroba, E. (2020). A review of *Listeria monocytogenes* from meat and meat products: Epidemiology, virulence factors, antimicrobial resistance and diagnosis. *Onderstepoort Journal of Veterinary Research*, 87(1). DOI:10.4102/ojvr.v87i1.1869.

11 Agentstvo po strategicheskemu planirovaniyu i reformam Respubliki Kazahstan Byuro nacional'noj statistiki. (2023). Statistika zdravoohraneniya i social'nogo obespecheniya, [in Russ].

12 Mezhgosudarstvennyisovetpostandartizacii,metrologiisertifikacii.(2014).Mezhgosudarstvennyj standart: Produkty pishchevye. Metody otbora prob dlya mikrobiologicheskikh ispytanij (GOST 31904-2012). Moskva: Standartinform, [in Russ].

13 Mezhgosudarstvennyi sovet po standartizacii, metrologii i sertifikacii. (2010). GOST 26669-85. Mezhgosudarstvennyi standart: Produkty pishchevye i vkusovye. Podgotovka prob dlya mikrobiologicheskogo analiza (GOST 26669-85). Moskva: Standartinform, [in Russ].

14 Mezhgosudarstvennyisovetpostandartizacii,metrologiisertifikacii.(2013).Mezhgosudarstvennyj standart: Produkty pishchevye. Metody vyyavleniya i opredeleniya kolichestva bakterij gruppy kishechnyh palochek (koliformnyh bakterij) (GOST 31747-2012). Moskva: Standartinform, [in Russ].

15 Mezhgosudarstvennyisovetpostandartizacii,metrologiisertifikacii.(2014).Mezhgosudarstvennyj standart: Produkty pishchevye. Metod vyyavleniya bakterij roda *Salmonella* (GOST 31659-2012). Moskva: Standartinform, [in Russ].

16 Mezhgosudarstvennyisovetpostandartizacii,metrologiisertifikacii.(2022).Mezhgosudarstvennyj standart: Produkty pishchevye. Metody vyyavleniya bakterij *Listeria monocytogenes* i drugih vidov *Listeria* (*Listeria spp.*) (GOST 32031-2022). Moskva: Standartinform, [in Russ].

17 Mezhgosudarstvennyisovetpostandartizacii,metrologiisertifikacii.(2013).Mezhgosudarstvennyj standart: Produkty pishchevye. Metody vyyavleniya i opredeleniya kolichestva koagulazopolozhitel'nyh stafilokokkov i *Staphylococcus aureus* (GOST 31746-2012). Moskva: Standartinform, [in Russ].

18 Mezhgosudarstvennyisovetpostandartizacii,metrologiisertifikacii.(2008).Mezhgosudarstvennyj standart: Produkty pishchevye. Metody kul'tivirovaniya mikroorganizmov (GOST 26670-91), Moskva: Standartinform. [in Russ].

19 Evrazijskaya ekonomicheskaya komissiya. (2011). Tekhnicheskij reglament Tamozhennogo soyuza TR TS 021/2011. O bezopasnosti pishchevoi produkcii. [in Russ].

20 Ngo, HHT, Nguyen-Thanh, L., Pham-Duc, P., Dang-Xuan, S., Le-Thi, H., Denis-Robichaud, J., Nguyen-Viet, H., Le, TT, Grace, D., Unger, F. (2021). Microbial contamination and associated risk factors in retailed pork from key value chains in Northern Vietnam. *International Journal of Food Microbiology*, 346, 109163. DOI: 10.1016/j.ijfoodmicro.2021.109163.

21 Veselá, H., Dorotíková, K., Dušková, M., Furmančíková, P., Šedo, O., Kameník, J. (2022). The pork meat or the environment of the production facility? The effect of individual technological steps on the bacterial contamination in cooked hams. *Microorganisms*, 10(6), 1106. DOI:10.3390/microorganisms10061106.

22 Lenchenko, E., Stepanov, D. (2020). Quantitative account and differential properties of pathogenic bacteria allocated from food raw materials. *Problems of Veterinary Sanitation Hygiene and Ecology*, 1(2), 228–235. DOI: 10.36871/vet.san.hyg.ecol.202002017.

23 Evrazijskaya ekonomicheskaya komissiya. (2013). Tekhnicheskii reglament Tamozhennogo soyuza TR TS 034/2013. O bezopasnosti myasa i myasnoj produkcii, [in Russ].

Herald of Science of S.Seifullin Kazakh Agrotechnical Research University: Veterinary Sciences.
– Astana: S. Seifullin Kazakh Agrotechnical Research University, 2024. – № 4 (008). – P. 43-50.
- ISSN 2958-5430, ISSN 2958-5449

doi.org/ 10.51452/kazatuvc.2024.4(008).1796
UDC581.192

Research article

Plants with radioprotective properties: current research and application prospects

Sergazy T. Dyusembayev¹ , Aynur T. Serikova¹ , Shyngys K. Suleimenov¹ ,
Zhaksylyk T. Serikov¹ , Zhanat Sh. Adilbekov² 

¹Shakarim University of Semey, Semey, Kazakhstan

²S.Seifullin Kazakh Agrotechnical Research University, Astana, Kazakhstan

Corresponding author: Shyngys K. Suleimenov: shyngys2203@mail.ru

Co-authors: (1: SD) sergazi_d@mail.ru; (2: AS) aiser_71@mail.ru;

(3: ShS) shyngys2203@mail.ru; (4: ZhS) axmedx93@mail.ru;

(5: ZhA) zh.adilbekov@kazatu.edu.kz

Received: 18-10-2024 **Accepted:** 26-12-2024 **Published:** 30-12-2024

Abstract

Background and Aim. The objective of this study is to investigate plants capable of reducing radionuclide accumulation in animals and to develop scientifically substantiated dietary feed supplements suitable for the conditions of the Semipalatinsk Nuclear Test Site (SNTS). Plants with radioprotective properties play a critical role in the context of the former SNTS.

Materials and Methods. The selection criteria for the plants included their ability to bind and remove radionuclides as well as their antioxidant activity. Key species studied included the fruits of wild apple, rosehip, rowan, and hawthorn. The selected plants were harvested, dried, and processed into powdered and extract forms. A series of experiments were conducted on livestock to test various feed supplements enriched with radioprotective plants and dietary compounds such as pectin and potassium ferrocyanide. These radioprotective agents were incorporated into animal feed at concentrations of 5–7% to evaluate their effectiveness. Radiometric monitoring in experimental areas was carried out using instruments such as the RKS-01-SOLO radiometer-dosimeter, XFC-AT 6130 dosimeter-radiometer, and RAMON-02 radon monitor. Highly sensitive dosimeters were used to determine radionuclide levels in plants.

Results. The use of radioprotective plants as feed additives reduced radionuclide content in meat by 15% compared to the control group. The application of radioprotective agents facilitated the accelerated elimination of radionuclides from animals' bodies. In the experimental group, the radionuclide excretion period was reduced to 15 days compared to the control group.

Conclusion. The study results demonstrate the effectiveness of radioprotective plants in reducing radioactive substances in animals. These plants exhibit the ability to bind and remove radionuclides from animals' bodies. Feed additives based on these plants enhance productivity, quality, and safety of livestock products in areas affected by the former SNTS.

Keywords: antioxidants; biologically active substances; radioprotectors; radiation exposure; plants; radiation.

Introduction

The radioprotective properties of plants are a focal point of contemporary scientific research, particularly due to the increasing risk of radiation exposure to the environment and animal health. Substances with radioprotective properties are crucial for radiation protection. The aim of this study is to identify and consolidate effective radioprotective substances and plants capable of reducing radionuclide

accumulation in animals, as well as to develop scientifically substantiated feed additives and diets for animals raised in radioactively contaminated areas, such as the former Semipalatinsk Nuclear Test Site (SNTS). The implementation of such additives for livestock in the SNTS region is an essential measure. Ensuring the quality and safety of meat, dairy, and other livestock products in the context of the former SNTS remains a pressing issue in veterinary science. Therefore, the adoption and use of feeds with radioprotective properties to reduce radionuclide accumulation in animals is imperative [1].

Numerous medicinal herbs exhibit radioprotective properties or the ability to bind and remove radionuclides from the body. Herbs such as licorice, eleuthero, ginseng, nettle, rockrose, plantain, chamomile, and horse chestnut help stabilize cellular membranes, reduce swelling, and suppress autoimmune processes. These herbs protect the liver and kidneys from damage, enhance tissue regeneration, stimulate hematopoiesis, support intestinal epithelium health, and normalize the functions of the immune system, liver, spleen, and thymus.

Herbs like meadowsweet and eleuthero contribute to the normalization of the sympathetic and parasympathetic nervous systems' tone [2].

A complex of endogenous biologically active compounds, including amines, thiols, and other antioxidants, boosts overall resistance to radiation and suppresses the excessive accumulation of radiation-induced oxidative products that are harmful to living cells. Herbs such as kelp and flaxseed act as sorbents, promoting the elimination of radionuclides from the body.

Based on the above, the objective of this study is to analyze current scientific data on the radioprotective properties of plants and substances capable of reducing radionuclide accumulation in animals [3].

Explosions at the SNTS resulted in the release of numerous radioactive elements into the environment, the most radiologically significant of which include iodine-131, cesium-137, strontium-90, plutonium isotopes (239, 240, 241), and americium-241 [4].

Once inside the body, radionuclides act as harmful agents, damaging cells and requiring removal through any available means. Prolonged exposure to even minimal doses can lead to cellular changes, causing genetic mutations, malignant tumors, and various metabolic disorders, including those affecting digestion, hematopoiesis, and other bodily functions.

Our previous studies (2012–2014) revealed that soil, vegetation, water sources, and livestock products in the SNTS area remain significantly contaminated, representing a large-scale radioecological problem with adverse consequences stemming from the former test site.

Currently, researchers have estimated that animal diets in the SNTS region lack 30–50% of essential vitamins, macroelements, and microelements. Consumption of food contaminated with radioactive substances can result in human health issues. Thus, it is crucial in this region to use foods enriched with high-quality protein, vitamins, macroelements, microelements, and substances with radioprotective properties. A distinctive feature of these radioprotective foods is their ability to normalize metabolic processes, address the root causes of metabolic disturbances, and reduce radionuclide accumulation in the body [5].

Destructive, antioxidant, and anti-inflammatory actions: herbs such as licorice, eleuthero, ginseng, nettle, rockrose, plantain, chamomile, and horse chestnut stabilize cell membranes, reduce swelling, and suppress autoimmune processes. Antimutagenic actions: phenolic compounds, beta-carotene, vitamin E, and other antioxidants contribute to the prevention of radiation-induced mutations. Vasodilatory effects: phenolic compounds help normalize vascular permeability and expand blood vessels. Detoxification: all these plants facilitate detoxification processes, aiding in the elimination of harmful substances from the body. Hepato- and nephroprotective actions: these herbs protect the liver and kidneys from damage. Regeneration and hematopoiesis: they promote tissue regeneration, stimulate hematopoiesis, support epithelial recovery in the colon, and normalize the functions of the immune system, liver, spleen, and thymus. Nervous system regulation: herbs such as meadowsweet and eleuthero help normalize the tone of the sympathetic and parasympathetic nervous systems. Enhancing non-specific resistance: a complex of endogenous biologically active compounds, including amines, thiols, and other antioxidants, increases general radiation resistance and suppresses the excessive accumulation of radiation-induced oxidative products harmful to living cells. Sorption properties: herbs such as kelp and flaxseed act as sorbents, promoting the excretion of radionuclides from the body [6].

The radiation situation in the former test site area remains complex. More than 30 years have passed since the closure of the SNTS, but the effects of nuclear explosions may persist for several decades [7].

Pectin is a gelatinous substance commonly found in fruit jams or jellies. During digestion, pectin binds with radionuclides and toxic heavy metals, forming insoluble salts that are excreted through the intestines. Additionally, low-molecular-weight pectin fractions can enter the bloodstream, where they form complexes with radionuclides and are subsequently excreted via urine. The most popular sources of pectin include apples, blackberries, rosehips, pears, and nettle. Products made from these fruits and plants are rich in biologically active substances with radioprotective properties. Naturally, berries such as blackcurrants, cherries, and apricots are rich in organic acids and other biologically active compounds that enhance their radioprotective effects [8, 9].

The development of a new type of dietary supplement is based on the inclusion of fishmeal rich in essential macro- and microelements, biologically necessary vitamins, radioprotective plants such as sea buckthorn, Jerusalem artichoke, rosehips, earth pear, and pectin substances. The combination of these beneficial properties provides radioprotective and preventive effects, opening avenues for the creation of scientifically substantiated food products [10].

In veterinary practice, the use of substances with radioprotective properties is crucial for ensuring high-quality and safe products by minimizing radioactive substances in animals, particularly sheep raised in the SNTS area. Certain products, such as soy, carrots, beets, turnips, cabbage, and mushrooms, are effective in removing radionuclides from the body. Additionally, increasing the consumption of potassium-rich foods such as eggplants, green peas, potatoes, tomatoes, and watermelons can help reduce radiocesium accumulation in critical organs [11, 12].

Wormwood (*Artemisia*): In Karaganda, Kazakh specialists developed an anticancer drug from local wormwood. This medication, known as arglabin, is widely used in oncology clinics across the Commonwealth of Independent States (CIS) and is patented in 11 nations. The drug enables the treatment and prevention of liver, lung, and breast tumors. Currently, arglabin has received 11 million international orders for cancer treatment and is in demand in Russia, Ukraine, Belarus, Tajikistan, and Uzbekistan. Domestic demand reaches 2 million units [13].

A diet rich in vitamins is essential, as numerous studies indicate that the body's demand for vitamins increases even under low doses of ionizing radiation. Moreover, certain vitamins can help the body tolerate better elevated levels of radiation [14]. The primary sources of vitamin E are unrefined vegetable oils, such as soybean, corn, sunflower, and rosehip oils. Carotenoids, which are precursors to vitamin A, are found in carrots, red peppers, peaches, apricots, sea buckthorn, rowanberries, rosehips, pumpkins, and ripe tomatoes. Vitamin C is particularly abundant in rosehips, blackcurrants, citrus fruits, green peas, zucchini, carrots, beets, radishes, cauliflower, dill, and other similar foods.

Previous research has highlighted that a specially designed feed mixture rich in vitamins B1, B2, B6, B12, and C, as well as mineral substances, exhibits radioprotective properties. It reduces fattening times by threefold, improves meat quality, increases animal live weight gain by up to 20%, and decreases the overall dry feed consumption by 6–14%. This next-generation feed mixture, developed for agricultural animals, combines biologically valuable, therapeutic, preventive, and specialized products to produce high-nutritional-value meat. This is particularly significant for populations residing near the SNTS [15].

A horse feed with radioprotective and anthelmintic properties utilizes wild red apples rich in minerals and vitamins. This feed mixture includes potassium ferrocyanide and the anthelmintic gel Brovermectin. Potassium ferrocyanide aids in the excretion of the radioactive substance cesium-137 from the horse's body, while Brovermectin eliminates parasitic helminths. The primary nutritional components of the feed are proteins, carbohydrates, lipids, vitamins, and other biologically active compounds. The high pectin content in the feed enhances its digestibility and contributes to its radioprotective properties [16].

A mineral-salt block has been created using feed-grade salt supplemented with wild apple, potassium ferrocyanide, and anthelmintic agents. Under the influence of these developed mineral-salt additives, the infection rates of sheep with strongyloidiasis and eimeriosis decreased from 49.8–73.75% to 3–7%. When using the mineral-salt blocks, the infection rates fell from 49.8–73.75% to 2–4%, and the levels of radioactive substances in the animals decreased by 12% [17].

At present, there is a pressing need to systematize and consolidate accumulated experience in this field, as well as to identify promising directions for further research and practical application.

Materials and Methods

Radiometric monitoring employs devices and instruments subject to annual state calibration, including the RKS-01-SOLO radiometer-dosimeter, the MKS-AT6130 dosimeter-radiometer, and the RAMON-2 radon meter. Research requires the establishment of control and measurement stations. Samples of vegetation and feeds are collected from these control stations according to the standards of the Republic of Kazakhstan, specifically following the sanitary rules outlined in SanPiN 6.01.001-97 *Unified Rules for Sampling Environmental Objects*, which are designed for veterinary, agrochemical, control-toxicological laboratories under the Ministry of Agriculture and other organizations.

In order to ensure reliable laboratory data on the contamination of vegetation and feeds, strict adherence to sampling rules is essential. Vegetation samples must be collected after ground surveys at designated plots measuring 1×1 m. The sample mass should be at least 300 g. The aboveground parts of plants are cut at a height of 3 cm. Sampling of plant-based feeds is conducted in accordance with GOST 27262-87 *Plant Feeds: Sampling Methods*. For plant-based feeds, organoleptic properties are evaluated.

The products must be free from foreign odors, tastes, inclusions, or other defects. When assessing the safety of vegetation, general toxicological properties, bacterial contamination (including the presence of conditionally pathogenic and pathogenic microorganisms), and the content of heavy metals, pesticides, nitrites, nitrates, mycotoxins, and other harmful impurities are determined.

The radiological safety of vegetation concerning cesium-137 and strontium-90 is assessed according to permissible levels of specific radionuclide activity established by radiation safety standards and sanitary rules GN 2.6.1.054-96, approved by the decree of the State Sanitary and Epidemiological Surveillance on April 19, 1996, No. 7.

Sample preparation: Data for each sample must be recorded in a field passport. In the laboratory, the vegetation is ground and homogenized. The extract is prepared as follows: weigh 2 g of the sample and place it in a round-bottom flask, add 20 mL of a 3% nitric acid solution, and mix the solution on a reciprocating shaker (LIOP LS 120) for 60 min. Filter the resulting suspension through paper filters. The filtrate is then diluted 50-fold with distilled water and analyzed using an optical emission spectrometer.

Stage 1: The effect of radioprotective feed additives on animal growth and development

Group 1: standard feed. Group 2: feed supplemented with sea buckthorn and Jerusalem artichoke. Group 3: feed supplemented with wormwood and potassium ferrocyanide. Experiment duration: 4 months. Measured parameters: live weight gain, radionuclide content in meat, and total feed consumption. According to the results of the experiment using feed supplemented with wormwood and potassium ferrocyanide (Group 3), the animals in this group demonstrated the greatest live weight gain and the lowest radionuclide content in meat. These findings were utilized to develop recommendations for feed additives with radioprotective properties.

Stage 2: Development and testing of a new phytopreparation

A new radioprotective agent was developed based on local plants, including rosehip, sea buckthorn, and wormwood, and its efficacy was tested on animals. For three months, animals in the experimental groups were administered the new phytopreparation, while the control group received no such treatment.

Results

Reduction in radionuclide accumulation

In the experimental groups fed with the phytopreparation based on sea buckthorn, Jerusalem artichoke, wormwood, and potassium ferrocyanide, a significant reduction in radionuclide accumulation (cesium-137, strontium-90) in tissues was recorded. In the group receiving wormwood and potassium ferrocyanide, the radionuclide content in meat was within permissible limits, whereas the control group showed a 15% exceedance of the norm (Table 1).

Table 1 – Radionuclide content in animal tissues (Bq/kg)

Study parameters	Control	Experimental
Average radionuclide content	25% above norm	10% above norm
Cesium-137	212.5 ± 0.05	205 ± 0.004
Strontium-90	0.004 ± 0.001	0.002 ± 0.0001

In the experimental groups the average radionuclide content after application of radioprotectors and potassium ferrocyanide decreased by 90%, and in the control group by 75% i.e. cesium-137 in the control group was found $212,5 \pm 0,05$ Bq/kg, and in the experimental group $205 \pm 0,004$ Bq/kg, respectively, strontium-90 - $0,004 \pm 0,001$ Bq/kg and $0,002 \pm 0,0001$ Bq/kg.

Weight gain and product quality improvement

In groups of animals fed with radioprotective supplements (sea buckthorn, Jerusalem artichoke, wormwood), a significant increase in live weight was observed: 20% in the group receiving wormwood and potassium ferrocyanide and 15% in the group fed sea buckthorn and Jerusalem artichoke. Moreover, the quality of meat products improved due to a reduction in radionuclide content and an increase in nutritional value (Table 2).

Table 2 – Diet and radionuclide content in meat

Animal group	Weight gain	Radionuclide content in meat	Radionuclide increase in meat
Group 1 (standard feed), average weight: 320 kg			
Cesium-137 Strontium-90	336 ± 0.04 kg	213 ± 0.001 Bq/kg 0.003 ± 0.001 Bq/kg	15% above norm
Group 2 (sea buckthorn, Jerusalem artichoke), average weight: 331 kg			
Cesium-137 Strontium-90	381 ± 0.05 kg	210 ± 0.001 Bq/kg 0.01 ± 0.001 Bq/kg	5% above norm
Group 3 (wormwood, potassium ferrocyanide), average weight: 330 kg			
Cesium-137 Strontium-90	396 ± 0.02 kg	180 ± 0.001 Bq/kg -	Normal

In the group not receiving radioprotective preparations, the amount of cesium-137 was found to be 213 ± 0.001 Bq/kg, and strontium-90 was 0.003 ± 0.001 Bq/kg, which is 15% higher than the norm.

In the second group, which received radioprotectors (Jerusalem artichoke and sea buckthorn) along with feed, the levels of cesium-137 and strontium-90 decreased by 95%.

In the third group, which received wormwood and ferrocyanide along with feed, no radioactive substances were detected. Adding sea buckthorn, Jerusalem artichoke, and potassium ferrocyanide to the diet significantly reduced the levels of radionuclides in the animals' meat. The lowest radionuclide content was observed in group 3.

Acceleration of radionuclide elimination

The research results indicate that plants with radioprotective properties may play a crucial role in radiation protection, particularly in regions affected by nuclear testing. Phytopreparations accelerated radionuclide elimination in animals, reducing the excretion period from 30 days in the control group to 15 days in the experimental group. Studies focused on the radioprotective properties of local plants in Kazakhstan, the analysis of their active compounds, and their impact on radiation protection. Feed additives based on these plants showed potential as environmentally friendly alternatives to synthetic radioprotectors, requiring further laboratory and field trials to confirm their safety and efficacy. A deeper analysis of plant-based radioprotectors is essential to understand their mechanisms and enhance their utilization. Systematizing data on these agents will contribute to the development of new methods and technologies for effective radiation protection. Research on plant-based radioprotectors opens promising opportunities for safer and more sustainable radiation protection strategies.

Discussion and Conclusion

The results of the conducted studies have shown high efficiency of radioprotective preparations and radioprotective feeds in reducing radiation contamination of animal organism and improving the quality of livestock products. Prospectivity of the use of plant radioprotectors: plants with radioprotective properties represent a promising alternative to synthetic radioprotectors, providing an environmentally friendly and potentially safe solution for protection from radiation exposure.

Comprehensive studies are needed to fully understand the mechanisms of action of plant radioprotectors, and systematization and generalization of available data on radioprotectors will help in the development of new methods and technologies, as well as in the practical application of radioprotectors.

Research suggests that plants with radioprotective properties may be key elements in a defense strategy against radiation exposure. The importance of further study of these plants, including their mechanisms of action, efficacy and safety, is undeniable. Comprehensive studies and systematization of data are needed to better understand the potential of plant radioprotectors. Thus, the study of plant radioprotectors is an important step towards the development of safer and more effective methods of protection against radiation exposure, with significant implications for science and practice in this field.

1. Plants with radioprotective properties, such as sea buckthorn, wormwood, Jerusalem artichoke, and rosehip, demonstrate the ability to bind and excrete radionuclides from the body while protecting cells from radiation-induced damage.

2. Fodder mixtures enriched with radioprotective plants and substances (e.g., pectin, potassium ferrocyanide) not only reduce radiation levels in animals but also enhance their productivity, improve the quality and safety of livestock products.

3. Radioprotective preparations derived from local plants accelerate radionuclide excretion and promote tissue regeneration, making them a promising solution for use in radiation-contaminated areas, such as the former Semipalatinsk nuclear test site.

4. The use of radioprotective additives shortens the fattening period, increases weight gain, and improves the biochemical parameters of blood - an essential factor for ensuring food security in regions with heightened radiation risk.

Authors' Contributions

SD and AS: Conceptualized and designed the study, conducted a comprehensive literature search, analyzed the gathered data and drafted the manuscript. ShS, ZhS and ZH: Conducted the final revision and proofreading of the manuscript. All authors have read, reviewed, and approved the final manuscript.

Acknowledgments

This research was conducted as part of the targeted funding program for scientific and scientific-technical programs for 2024-2026 (Ministry of Foreign Affairs, Republic of Kazakhstan), Project ID: BR22885795, Project Title: "Improving food product safety."

References

- 1 Роева, НН, Зайцева, ИА. (2023). Аналитический контроль содержания радионуклидов в кормах. *Управление качества*, 2. DOI: 10.33920/pro-01-2302-01.
- 2 Lee, T., Johnke, R., Allison, R., O'Brien, K., Dobbs, L. (2005). Radioprotective potential of ginseng. *Mutagenesis*, 20(4), 237-243. DOI:10.1093/MUTAGE/GEI041.
- 3 Garder, K., Skulberg, O. (1965). Radionuclide accumulation by *Anodonta piscinalis* Nilsson (Lamellibranchiata) in a continuous flow system. *Hydrobiologia*, 26, 151-169. DOI:10.1007/BF00142263.
- 4 Nakamura, T., Masuda, S., Kuchiki, A., Maruyama, A. (2020). Effects of radioactive contamination from the Semipalatinsk nuclear test site on behavior related to food choices: A case study of Kazakhstan. *Journal of Disaster Research*, 15, 991-1010. DOI:10.20965/jdr.2020.p0991.
- 5 Krasnopoyrova, M., Gorlachev, I., Kharkin, P., Dyussebayeva, M., Lukashenko, S. (2023). Study of radionuclide composition of the drinking water potential sources at the Semipalatinsk nuclear test site. *RSC Advances*, 13, 29354-29362. DOI:10.1039/d3ra05808e.
- 6 Mueed, A., Shibli, S., Korma, S., Madjirebaye, P., Esatbeyoglu, T., Deng, Z. (2022). Flaxseed bioactive compounds: Chemical composition, functional properties, food applications and health benefits-related gut microbes. *Foods*, 11. DOI:10.3390/foods11203307.

7 Смагулова, ГМ, Тулеуова, БТ, Стамбулов, СБ, Кенжегали, ЖМ. (2023). Семей полигоны: тарихы, ядролық сынақтар және оның зардаптары. *Вестник Карагандинского университета. Серия История. Философия*, 3(111), 113-120. DOI:10.31489/2023hph3/113-120.

8 Jandosov, J., Alavijeh, M., Sultakhan, S., Baimenov, A., Bernardo, M., Sakipova, Z., Azat, S., Lyubchuk, S., Zhylybayeva, N., Naurzbayeva, G., Mansurov, Z., Mikhalovsky, S., Berillo, D. (2022). Activated carbon/pectin composite enterosorbent for human protection from intoxication with xenobiotics Pb (II) and sodium diclofenac. *Molecules*, 27. DOI:10.3390/molecules27072296.

9 Кокоев, МН. (2023). Новый способ измельчения сырья для производства пектина. *Вестник российской академии естественных наук*, 23(3), 23-28. DOI:10.52531/1682-1696-2023-23-3-23-28.

10 Dowlath, M., Karuppannan, S., Sinha, P., Dowlath, N., Arunachalam, K., Ravindran, B., Chang, S., Nguyen-Tri, P., Nguyen, D. (2021). Effects of radiation and role of plants in radioprotection: A critical review. *The Science of the Total Environment*, 779, 146431. DOI: 10.1016/J.SCITOTENV.2021.146431.

11 Тюрин, ВГ, Семенов, ВГ, Вагин, КН, Гайнутдинов, ТР, Низамов, РН, Галлямова, МЮ, Авыллов, ЧК. (2023). Санитарно-гигиенические показатели продуктов убоя при радиационных поражениях животных на фоне применения радиопротектора на основе веществ микробного происхождения. *Российский журнал «Проблемы ветеринарной санитарии, гигиены и экологии»*, 1(45). DOI: 10.36871/vet.san.hyг.ecol.202301010.

12 Panitskiy, A., Bazarbaeva, A., Baigazy, S., Polivkina, Y., Alexandrovich, I., Abisheva, M. (2023). Bioaccumulation of radionuclides in hoofed animals inhabiting the Semipalatinsk Test Site. *PLOS ONE*, 18. DOI: 10.1371/journal.pone.0294632.

13 Неверова, НА, Жабаева, АН, Левчук, АА, Бабкин, ВА, Бейсенбаев, АР, Ларина, ЛИ, Сапожников, АН, Адекенов, СМ. (2019). Исследование физико-химических свойств механообработанного арглабина и его механокомпозигов на основе арабиногалактана. *Химия растительного сырья*, 4, 106-107. DOI:10.14258/jcprm.2019046259.

14 Ponnampalam, E., Kiani, A., Santhiravel, S., Holman, B., Lauridsen, C., Dunshea, F. (2022). The importance of dietary antioxidants on oxidative stress, meat and milk production, and their preservative aspects in farm animals: Antioxidant action, animal health, and product quality-Invited review. *Animals: An Open Access Journal from MDPI*, 12, 21-22. DOI:10.3390/ani12233279.

15 Obeid, R., Heil, S., Verhoeven, M., Heuvel, E., Groot, L., Eussen, S. (2019). Vitamin B12 intake from animal foods, biomarkers, and health aspects. *Frontiers in Nutrition*, 6. DOI:10.3389/fnut.2019.00093.

16 Bampidis, V., Azimonti, G., Bastos, M., Christensen, H., Dusemund, B., Durjava, M., Kouba, M., López-Alonso, M., Puente, S., Marcon, F., Mayo, B., Pechová, A., Petkova, M., Ramos, F., Sanz, Y., Villa, R., Woutersen, R., Aquilina, G., Bories, G., Nebbia, C., Gropp, J., Galobart, J., Vettori, M., Anguita, M., Casanova, J., Amaduzzi, A., Innocenti, M. (2023). Safety and efficacy of feed additives consisting of sodium ferrocyanide and potassium ferrocyanide for all animal species (Eusalt a.i.s.b.l.). *EFSA Journal*, 21, 7960. DOI: 10.2903/j.efsa.2023.7960.

17 Pareja-Carrera, J., Martínez-Haro, M., Mateo, R., Rodríguez-Estival, J. (2020). Effect of mineral supplementation on lead bioavailability and toxicity biomarkers in sheep exposed to mining pollution. *Environmental Research*, 110364. DOI: 10.1016/j.envres.2020.110364.

References

1 Roeva, NN, Zaytseva, IA. (2023). Analiticheskii kontrol' soderzhaniya radionuklidov v kormakh. *Upravlenie kachestva*, 2. DOI:10.33920/pro-01-2302-01.

2 Lee, T., Johnke, R., Allison, R., O'Brien, K., Dobbs, L. (2005). Radioprotective potential of ginseng. *Mutagenesis*, 20(4), 237-243. DOI:10.1093/MUTAGE/GEI041.

3 Garder, K., Skulberg, O. (1965). Radionuclide accumulation by Anodonta piscinalis Nilsson (Lamellibranchiata) in a continuous flow system. *Hydrobiologia*, 26, 151-169. DOI:10.1007/BF00142263.

4 Nakamura, T., Masuda, S., Kuchiki, A., Maruyama, A. (2020). Effects of radioactive contamination from the Semipalatinsk nuclear test site on behavior related to food choices: A case study of Kazakhstan. *Journal of Disaster Research*, 15, 991-1010. DOI:10.20965/jdr.2020.p0991.

- 5 Krasnopyorova, M., Gorlachev, I., Kharkin, P., Dyussebayeva, M., Lukashenko, S. (2023). Study of radionuclide composition of the drinking water potential sources at the Semipalatinsk nuclear test site. *RSC Advances*, 13, 29354-29362. DOI:10.1039/d3ra05808e.
- 6 Mueed, A., Shibli, S., Korma, S., Madjirebaye, P., Esatbeyoglu, T., Deng, Z. (2022). Flaxseed bioactive compounds: Chemical composition, functional properties, food applications and health benefits-related gut microbes. *Foods*, 11. DOI:10.3390/foods11203307.
- 7 Ganiev, GM, Tursynova, BT, Sarsenbaev, SB, Kenzhina, ZM. (2023). Semipalatinskii ispytatel'nyy poligon: istoriya, yadernyye ispytaniya i posledstviya. *Vestnik Karagandinskogo universiteta. Seriya Istoriya. Filosofiya*, 3(111), 113-120. DOI:10.31489/2023hph3/113-120.
- 8 Jandosov, J., Alavijeh, M., Sultakhan, S., Baimenov, A., Bernardo, M., Sakipova, Z., Azat, S., Lyubchik, S., Zhylybayeva, N., Naurzabayeva, G., Mansurov, Z., Mikhalovsky, S., Berillo, D. (2022). Activated carbon/pectin composite enterosorbent for human protection from intoxication with xenobiotics Pb (II) and sodium diclofenac. *Molecules*, 27. DOI:10.3390/molecules27072296.
- 9 Kokoev, MN. (2023). Novyi sposob izmel'cheniya syrya dlya proizvodstva pektina. *Nauchnaya stat'ya*, 23(3), 23-28. DOI:10.52531/1682-1696-2023-23-3-23-28.
- 10 Dowlath, M., Karuppanan, S., Sinha, P., Dowlath, N., Arunachalam, K., Ravindran, B., Chang, S., Nguyen-Tri, P., Nguyen, D. (2021). Effects of radiation and role of plants in radioprotection: A critical review. *The Science of the Total Environment*, 779, 146431. DOI: 10.1016/J.SCITOTENV.2021.146431.
- 11 Tyurin, V., Semenov, V., Vagin, K., Gaynutdinov, T., Nizamov, R., Gallyamova, M., Avylov, C. (2023). Sanitarno-gigienicheskie pokazateli produktov uboja pri radiacionnyh porazheniyah zhivotnyh na fone primeneniya radioprotektora na osnove veshhestv mikrobnogo proishozhdeniya. *Rossiiskii zhurnal «Problemy veterinarnoy sanitarii, gigieny i ekologii»*, 1(45). DOI: 10.36871/vet.san.hyg.ecol.202301010.
- 12 Panitskiy, A., Bazarbaeva, A., Baigazy, S., Polivkina, Y., Alexandrovich, I., Abisheva, M. (2023). Bioaccumulation of radionuclides in hoofed animals inhabiting the Semipalatinsk Test Site. *PLOS ONE*, 18. DOI: 10.1371/journal.pone.0294632.
- 13 Neverova, NA, Zhabaeva, AN, Levchuk, AA, Babkin, VA, Bejsenbaev, AR, Larina, LI, Sapozhnikov, AN, Adekenov, SM. (2019). Issledovanie fiziko-himicheskikh svoystv mehanoobrabotannogo arglabina i ego mehanokompozitov na osnove arabinogalaktana. *Himija rastitel'nogo syr'ya*, 4, 106-107. DOI:10.14258/jcprm.2019046259.
- 14 Ponnampalam, E., Kiani, A., Santhiravel, S., Holman, B., Lauridsen, C., Dunshea, F. (2022). The importance of dietary antioxidants on oxidative stress, meat and milk production, and their preservative aspects in farm animals: Antioxidant action, animal health, and product quality-Invited review. *Animals: An Open Access Journal from MDPI*, 12, 21-22. DOI:10.3390/ani12233279.
- 15 Obeid, R., Heil, S., Verhoeven, M., Heuvel, E., Groot, L., Eussen, S. (2019). Vitamin B12 intake from animal foods, biomarkers, and health aspects. *Frontiers in Nutrition*, 6. DOI:10.3389/fnut.2019.00093.
- 16 Bampidis, V., Azimonti, G., Bastos, M., Christensen, H., Dusemund, B., Durjava, M., Kouba, M., López-Alonso, M., Puente, S., Marcon, F., Mayo, B., Pechová, A., Petkova, M., Ramos, F., Sanz, Y., Villa, R., Woutersen, R., Aquilina, G., Bories, G., Nebbia, C., Gropp, J., Galobart, J., Vettori, M., Anguita, M., Casanova, J., Amaduzzi, A., Innocenti, M. (2023). Safety and efficacy of feed additives consisting of sodium ferrocyanide and potassium ferrocyanide for all animal species (Eusalt a.i.s.b.l.). *EFSA Journal*, 21, 7960. DOI: 10.2903/j.efsa.2023.7960.
- 17 Pareja-Carrera, J., Martínez-Haro, M., Mateo, R., Rodríguez-Estival, J. (2020). Effect of mineral supplementation on lead bioavailability and toxicity biomarkers in sheep exposed to mining pollution. *Environmental Research*, 110364. DOI: 10.1016/j.envres.2020.110364.









Herald of Science of S.Seifullin Kazakh Agrotechnical Research University: Veterinary Sciences.
– Astana: S. Seifullin Kazakh Agrotechnical Research University, 2024. – № 4 (008). – P. 51-58.
- ISSN 2958-5430, ISSN 2958-5449

doi.org/ 10.51452/kazatuvc.2024.4(008).1811

UDC 579.23'314:616-097.155.37

Research article

Antigenic Activity of Various *Rhodococcus equi* Strain Plasmids

Gulnaz D. Ilgekbayeva , Yerken.I. Kassymov , Bauyrzhan. K. Otarbayev ,
Makpal Z. ZaniLabdin , Serikzhan Kurman , Nurali Matikhan , Ulan Zholdas ,
Bayan A. Valieva 

Kazakh National Agrarian Research University, Almaty, Kazakhstan

Corresponding author: Gulnaz D. Ilgekbayeva: gulnaz66@mail.ru

Co-authors: (1: YK) yerken.kassymov@mail.ru; (2: BO) bauken_68@mail.ru;
(3: MZ) m.zanilabdin@mail.ru; (4: CK) kurmanserikzhan79@gmail.com;
(5: NM) nurik200886@mail.ru; (6: UZ) ulanzholdass@gmail.com; (7: BV) v.ba.yan@mail.ru

Received: 02-12-2024 **Accepted:** 20-12-2024 **Published:** 30-12-2024

Abstract

Background and Aim. *Rhodococcus equi* (*R. equi*) infection is a fatal cause of equine *rhodococcosis*. Infections have also been reported in other species and humans. This study evaluated the immune response of *R. equi* in rabbits.

Materials and Methods. Different strains of *R. equi* (with and without plasmids) were administered to rabbits. Blood samples were collected from the ear veins of the rabbits at 7, 14, 21, and 28 days after the initial administration of the antigen, and the serum was examined for specific antibodies against *R. equi* plasmid antigens using the complement fixation reaction.

Results. The minimum antibody level was recorded on day 7. The average antibody level throughout the study period was slightly above the median, indicating a small asymmetry toward higher values. By day 28, the antibody level had reached 75% of values ≤ 4.25 .

Conclusion. All strains of *R. equi* exhibited antigenic properties to varying degrees. The antibody level upon plasmid pVapN administration was higher than that of the other cases.

Keywords: *Rhodococcus equi*; plasmid pVapB; plasmid pVapA; plasmid pVapN; antigenic activity.

Introduction

Rhodococcus equi (*R. equi*) is a pleomorphic Gram – positive *coccobacillus bacterium* commonly found in the gastrointestinal tract of horses. *R. equi* is recognized as an important pathogen based on reports of 3% mortality in foals, which is widespread globally [1, 2, 3]. *R. equi* is a primary lung pathogen in foals aged 6 months with high mortality rates [4, 5]. Pneumonia caused by *R. equi* can also occur in adult horses, especially in immunocompromised individuals, who can develop systemic infections [6].

In infected animals, chronic and purulent bronchopneumonia is typically observed and is associated with high mortality, especially in foals that have not been subjected to specific antibacterial therapy [3, 7]. Numerous studies using virulent strains of *R. equi* isolated from horses in different geographical regions worldwide have demonstrated the diversity of plasmid sizes and the predominance of different types of plasmids [8, 9, 10, 11, 12].

It has been established that only virulent strains of *R. equi*, which express the virulence-associated protein (VapA) of 15-17 kDa and possess a large virulence plasmid of 85-90 kDa containing the VapA gene, are pathogenic to horses. To date, 12 types of plasmids have been identified in VapA-positive strains from horses [11, 13, 14, 15]. VapA and VapB, which are closely related at the amino acid level, are located on the cell surface, and their expression is regulated by temperature and pH [16].

In studies on the source of intermediate virulence, VapB-positive strains have been isolated from the submandibular lymph nodes of pigs; these isolates were found to be of intermediate virulence in mice and contained one of five large plasmids sized 79±95 kb [17]. Virulent strains of *Rhodococcus equi* expressing the virulence-associated protein of 15–17 kDa (VapA) and possessing a large virulence plasmid (pVAPA) of 85–90 kb, containing the vapA gene are pathogenic to horses. Over the last two decades, after pVAPA, two types of host-associated virulence plasmids of *R. equi* have been discovered: the circular plasmid pVAPB, associated with isolates from pigs in 1995, and the recently identified linear plasmid pVAPN, linked to isolates from cattle and goats.

Molecular epidemiological studies of *R. equi* infections in foals on equine breeding farms in Japan and many countries around the world have been conducted over the past three decades, and epidemiological studies using the digestion schemes of plasmid DNA from virulent isolates with restriction enzymes have shown 14 different subtypes of pVAPA and their geographical preferences [18, 19, 20].

Virulent strains of *R. equi* isolated from sick foals and horse breeding environments exhibit a uniform plasmid pattern [21]. Certain types of plasmids are characteristic of specific geographical regions, which can be used in epidemiological studies [10, 11].

The aim of our research was to study the antigenic activity of various plasmids of the *R. equi* strain.

Materials and Methods

To study the antigenic activity of the plasmids of the *R. equi* strain, rabbits were immunized. We used the following strains, provided by Kitasato University (Japan):

- 1) The *Yokkaichi P* strain contains no plasmids.
- 2) S4 cells containing the plasmid pVapB.
- 3) ATCC33701 cells carrying the plasmid pVapA.
- 4) Yokkaichi strain containing the plasmid pVapN.
- 5) The *R. equi* strain was isolated by us in 2023 [22].

Cultivation of *Rhodococcus equi* Suspensions: For inoculation, a 2-day culture of the strain in tubes was washed with saline solution to achieve a microbial cell concentration of 2–5 billion cells in 1.0 cm³, according to the optical turbidity standard. This suspension was used to inoculate Tartakovsky flasks with NANAT medium [23]. The inoculated flasks were supplemented with 4.0–5.0 cm³ of suspension and placed in a thermostatic incubator at 37 °C for 2 days. After 2 days, the cultured material was visually inspected for purity and typical growth characteristics.

Preparation of the antigens: The culture was washed with a sterile 0.5% phenolized saline solution, pH 7.0–7.2, using 25–30 cm³ per flask. The resulting suspension, containing approximately 20–30 billion microbial cells in 1.0 cm³ according to the turbidity standard, was filtered through a double-layered gauze filter into bottles. The purity of the gram-stained smears. The mixture was then heated in a water bath at +70 °C for 60 min.

After cooling, the heated suspension was stored in a refrigerator at +2 °– 4 °C, where it was also checked for purity and sterility.

Sterility determination: From the antigen stock, the samples were inoculated onto MPA and NANAT media. The cultures were incubated in a thermostat for 3 days at a temperature of 37 °C–38 °C. The cultures remained sterile throughout the observation period.

In each group, three rabbits weighing 2.0–2.5 kg were selected and subcutaneously injected with the antigen at a dose of 2 cm³ twice, with an interval of 14 days. Blood samples were collected from the ear veins of the rabbits at 7, 14, 21, and 28 days following the first antigen administration, and the serum was tested for specific antibodies against the plasmid antigens of *Rhodococcus equi* using the prolonged complement fixation test (PCFT).

PCFT was performed under cold conditions in tubes with 0.2 cm³ volumes of serum, antigen, and complement, with the hemolytic system in a titrated working dose. Each antigen was titrated with the corresponding rabbit serum using the checkerboard method in PCFT. Serum samples were collected from rabbits on days 0, 7, 14, 21, and 28 of the experiment.

Statistical Analysis: Data analysis was performed using R-Studio software, applying the non-parametric Kruskal–Wallis test. Statistical significance was assessed at a threshold level of $P < 0.05$ [24].

The experiments and methodology used to conduct research on laboratory animals comply with the requirements of biological safety and the ethical principles of animal experimentation in the European Convention for the Protection of Vertebrate Animals used for Experimental and Other Scientific Purposes (Conclusion of the Bioethics Commission of the Kazakh National Agrarian Research University dated November 2, 2022).

Results

The checkerboard titration of the antigen and rabbit serum before antigen administration yielded a negative result. The results of the studies, starting from the seventh day of the experiment, are presented in Table 1.

Table 1 – Antibody Levels in Rabbits Administered Different Strains (with no plasmids, plasmids) of *R. equi*

Days of the Experiment	Group number-Average antibody titer				
	1 - no plasmids	2 - pVapB	3 - pVapA	4 - pVapN	5 - <i>R. equi</i>
7	-	1:5	1:5	1:5	1:5
14	-	1:10	1:10	1:10	1:10
21	1:10	1:20	1:20	1:40	1:20
28	1:20	1:80	1:40	1:80	1:40

As can be seen from the table, when the antigen from the strain without plasmids was injected into the group of rabbits on days 7 and 14, no antibodies were detected in the PCFT, whereas in the groups of rabbits injected with pVapB, VapA, pVapN, and *R. equi*, the antibody titer was within 1:5 and 1:10, respectively.

From day 21 in the first group, antibody levels began to appear at 1:10 and peaked at 1:20 on day 28. On day 28 of the experiment, a significant increase in antibody titer was observed in rabbits in groups 2 and 4, i.e., when antigen from the strain carrying the pVapB and pVapN plasmids was administered.

Because of the studies in the first group, rabbit serum antigen had the highest titer at dilutions of 1:50, 1:100, and 1:150. In serum samples positive for the strain without plasmids, the indicator was 1:10. With a decrease or increase in antigen dilutions, a decrease or absence of antibody levels is observed. An increase or decrease in antibody titers is directly dependent on the dilution of the antigen.

The absence of anticomplementary properties was demonstrated in the reactions of antigen dilutions with physiological solutions.

The antigen dose with a high antibody response in the reactions was taken as the equivalent zone. In addition to the equivalent zone, a decrease in the reaction results was observed at other antigen dilutions.

Because of the reactions, the dilution of the antigen that gave the highest antibody titer in the serum was taken as the antigen unit (AU). This AU is the most effective antigen dilution.

It was established that 1 AU for an antigen without plasmids is 1:150, with a working dose of the antigen of 1:150 equivalent to 1 AU. Higher than 1 AU resulted in self-limiting reactions.

In the checkerboard titration of rabbit serum treated with the antigen from the plasmid pVapB, the following results were obtained: 1 AU was 1:250, and the working dose of the antigen was 1:200, which corresponds to 1.25 AU.

For the titration of rabbit sera and the antigen from the plasmid pVapA, 1 AU was 1:300, and the working dose was 1:250, resulting in 1.2 AU.

Regarding pVapN, 1 AU was 1:200, while the working dose was 1:250 (1.3 AU). The 1 AU of the antigen from the *R. equi* strain was 1:300, whereas the working dose was 1:250 (1.2 AU).

The minimum antibody level was recorded on day 7. 25% of the values were ≤ 2.0 , which coincides with the level on day 14. The median value indicates the central tendency of the data. The average antibody level over the entire period was slightly above the median, indicating a small asymmetry toward higher values. By day 28, the antibody level had reached 75% of values ≤ 4.25 .

A P value of >0.05 indicates the absence of significant differences between groups. Variations in antibody levels are more likely related to variability than to the action of the plasmids.

Plasmids did not have a significant effect on antibody levels. However, the antibody level upon the administration of the plasmid pVapN (3.0) was somewhat higher than that in the other cases.

A clear increase in antibody titers was observed in rabbits with increasing time after the administration of the strains. This indicates the formation of an immune response. The dynamics of antibody titers showed a significant increase in antibody levels as the observation time increased, indicating the activation of the immune response. The differences between days were statistically significant ($P < 0.01$).

Analysis of the impact of plasmids on antibody production indicated that the differences between groups with various plasmids and those without plasmids were not statistically significant ($P = 0.62$). Nevertheless, plasmid pVapN showed a slight trend toward increased antibody levels.

The values of the antibody titers exhibited a slight asymmetry toward an increase, suggesting a rise in the antibody levels in later days of the experiment.

Discussion and Conclusion

Pathogenic strains are classified as virulent or intermediate virulent based on the presence of plasmid genes encoding virulence-associated protein A (VapA) or B (VapB), respectively [25]. Strains lacking VapA or VapB are classified as avirulent. The highly conserved gene encoding the conjugative transfer protein, traA, is present in strains carrying the virulence plasmid.

The new linear virulence plasmid *R. equi* (pVapN) has been characterized in isolates from cattle [26] and identified in a single isolate from dogs [27], but it has not been described in other species. Epidemiological data indicate that strains carrying pVAPN are pathogenic to ruminants [19].

VapN-positive *R. equi* has been isolated from cattle in Japan. Although *R. equi* is generally considered to have low pathogenicity in cattle, the influence of certain predisposing factors can lead to widespread infection [28].

VapA is an important antigen involved in the humoral protective immune response to *R. equi* infections caused by virulent strains in foals [29].

In this study, specific *R. equi* strains that are known to be pathogenic in various animal species were used. In our experience with rabbits, they elicited an immune response characterized by a clear increase in antibody titers over time following the administration of the strains.

All strains of *R. equi* (with and without plasmids, as well as the strain itself) exhibited antigenic properties to varying degrees. The antigen from the *R. equi* strain without plasmids showed the lowest level of activity. The antibody level upon administration of the plasmid pVapN was somewhat higher than that in the other cases.

Statistical analysis showed that the differences between the means at each stage of the experiment were statistically significant ($P < 0.01$), confirming a progressive increase in the immune response and greater variability at later stages of follow-up.

“Antigenic unit” was used as a measure of antigenic activity in serological reactions. The highest antigen titer corresponding to the highest value of positive blood serum with antibodies of the corresponding type was used as the antigenic unit. The value of the antigenic unit is not affected by the activity of the specific blood serum used to determine it; thus, the antigenic unit can be determined with any positive blood serum by titration using the checkerboard method.

The results may help obtain hyperimmune (positive) serum for the development of diagnostic kits for *rhodococcosis*. This study should be considered as a pilot study for the further development of *R. equi* bioproducts in horses and goats.

Authors' Contributions

GI, EK, and BO: Conceptualized and designed the study, conducted a comprehensive literature search, analyzed the collected data, and prepared the manuscript. MZ, SK, and NM: Prepared materials for hyperimmunization's and assessed antigen activity. UZ and BV: conducted the hyperimmunization of rabbits. All authors read, reviewed, and approved the final manuscript.

Acknowledgments

This work was supported by grant funding from the Science Committee of the Ministry of Science and Higher Education of the Republic of Kazakhstan for 2023-2025, grant AP19680565 “Creation of an

experimental infection model in goats using virulent *Rhodococcus equi* containing pVAPN as a model for *Rhodococcus* infection in foals.”

We express our gratitude to Sh. Takai (Kitasato University, Japan) for kindly providing the plasmids of the *Rhodococcus equi* strain, and to Yerlan Durbayev (Department of Plant Protection, Kazakh National Agrarian Research University, Almaty, Kazakhstan) for conducting the statistical analysis of the obtained data.

References

- 1 Takai, S., Mizuno, Y., Suzuki, Y., Sasaki, Y., Kakuda, T., Kirikae, T. (2024). *Rhodococcus equi* infections in humans: an emerging zoonotic pathogen. *Nihon Saikingaku Zasshi*, 79(1), 15-24. DOI: 10.3412/jsb.79.15.
- 2 Val-Calvo, J., Darcy, J., Gibbons, J., Creighton, A., Egan, C., Buckley, T. et al. (2022). International Spread of Multidrug-Resistant *Rhodococcus equi*. *Emerging infectious diseases*, 28(9), 1899-903.
- 3 Rakowska, A., Cywinska, A., Witkowski, L. (2020). Current Trends in Understanding and Managing Equine Rhodococcosis. *Animals*, 10(10), 1910. DOI: 10.3390/ani10101910.
- 4 Lankenfeld, A., Weber, C., Rohn, K., Venner, M. (2021). Kinetics of serum amyloid A during the treatment period of foals with pneumonia. *Pferdeheilkunde – Equine Medicine*, 37.
- 5 Arnold-Lehna, D., Venner, M., Berghaus, LJ, Berghaus, R., Giguère, S. (2020). Changing policy to treat foals with *Rhodococcus equi* pneumonia in the later course of disease decreases antimicrobial usage without increasing mortality rate. *Equine veterinary journal*, 52(4), 531-7.
- 6 Bordin, AI, Huber, L., Sanz, MG, Cohen, ND. (2022). *Rhodococcus equi* foal pneumonia: Update on epidemiology, immunity, treatment and prevention. *Equine Vet J*, 54(3), 481-94.
- 7 Cohen, ND. (2014). *Rhodococcus equi* foal pneumonia. *Vet Clin North Am Equine Pract*, 30, 609–622.
- 8 Venner, M., Meyer-Hamme, B., Verspohl, J., Hatori, F., Shimizu, N., Sasaki, Y. et al. (2007). Genotypic characterization of VapA positive *Rhodococcus equi* in foals with pulmonary affection and their soil environment on a warmblood horse breeding farm in Germany. *Res Vet Sci.*, 83, 311–317.
- 9 Son, WG, Lee, DS, Yamatoda, N., Hatori, F., Shimizu, N., Kakuda, T. et al. (2006). Molecular typing of VapA-positive *Rhodococcus equi* isolates from Jeju native horses, Korea. *J Vet Med Sci.*, 68, 249–253.
- 10 Duquesne, F., Hébert, L., Sévin, C., Breuil, MF, Tapprest, J., Laugier, C. et al. (2010). Analysis of plasmid diversity in 96 *Rhodococcus equi* strains isolated in Normandy (France) and sequencing of the 87-kb type I virulence plasmid. *FEMS Microbiol Lett.*, 311, 76–81.
- 11 Takai, S., Chaffin, MK, Cohen, ND, Hara, M., Nakamura, M., Kakuda, T. et al. (2001). Prevalence of virulent *Rhodococcus equi* in soil from 5 R. equi-endemic horse-breeding farms and restriction fragment length polymorphisms of virulence plasmids in isolates from soil and infected foals in Texas. *J Vet Diagn Invest.*, 13, 489–494.
- 12 Sangal, V., Goodfellow, M., Jones, AL, Sutcliffe, IC. (2022). A stable home for an equine pathogen: valid publication of the binomial *Prescottella equi* gen. nov., comb. nov., and reclassification of four rhodococcal species into the genus *Prescottella*. *Int J Syst Evol Microbiol.*, 72(9).
- 13 Takai, S., Sekizaki, T., Ozawa, T., Sugawara, T., Watanabe, Y., Tsubaki, S. (1991). Association between a large plasmid and 15- to 17-kilodalton antigens in virulent *Rhodococcus equi*. *Infect Immun.*, 59, 4056–4060.
- 14 Takai, S., Shoda, M., Sasaki, Y., Tsubaki, S., Fortier, G., Pronost, S. et al. (1999). Restriction fragment length polymorphism of virulence plasmids in *Rhodococcus equi*. *J Clin Microbiol.*, 37, 3417–3420.
- 15 Takai, S., Son, WG, Lee, DS, Madarame, H., Seki, I., Yamatoda, N. et al. (2003). *Rhodococcus equi* virulence plasmids recovered from horses and their environment in Jeju, Korea: 90-kb type II and a new variant, 90-kb type. *V. J Vet Med Sci.*, 65, 1313–1317.
- 16 Takai, S., Suzuki, Y., Sasaki, Y., Kakuda, T., Ribeiro, MG, Makrai, L., Witkowski, L., Cohen, N., Sekizaki, T. (2023). Short review: Geographical distribution of equine-associated pVAPA plasmids in *Rhodococcus equi* in the world. *Vet Microbiol.*, 287, 109919. DOI: 10.1016/j.vetmic.2023.109919.

- 17 Takai, S., Anzai, T., Fujita, Y., Akita, O., Shoda, M., Tsubaki, S., Wada, R. (2000). Pathogenicity of *Rhodococcus equi* expressing a virulence-associated 20 kDa protein (VapB) in foals. *Veterinary Microbiology*, 76: 1, 71-80.
- 18 Ford, J., Santos-Aberturas, J., Hems, E., Sallmen, J., Bögeholz, L., Polturak, G., Osbourn, A., Wright, J., Rodnina, M., Vereecke, D., Francis, I., Truman, A. (2024). Discovery of lydiamycin A biosynthetic gene cluster in the plant pathogen *Rhodococcus fascians* guides structural revision and identification of molecular target. *BioRxiv*, DOI: 10.1101/2024.11.13.623425.
- 19 Suzuki, Y., Takahashi, K., Ishitsuka, T., Sugiyama, M., Sasaki, Y., Kakuda, T., Takai, S., Naito, I., Kohara, J. (2024). Experimental infection of goats with pVAPN-harboring *Rhodococcus equi* causes latent infection in the lymph nodes. *Vet Microbiol.*, 295, 110132. DOI: 10.1016/j.vetmic.2024.110132.
- 20 Suzuki, Y., Takahashi, K., Takase, F., Sawada, N., Nakao, S., Toda, A., Sasaki, Y., Kakuda, T., Takai, S. (2020). Serological epidemiological surveillance for vapN-harboring *Rhodococcus equi* infection in goats. *Comp Immunol Microbiol Infect Dis.*, 73, 101540. DOI: 10.1016/j.cimid.2020.101540.
- 21 Kalinowski, M., Grądzki, Z., Jarosz, Ł., Kato, K., Hieda, Y., Kakuda, T., Takai, S. (2016). Plasmid Profiles of Virulent *Rhodococcus equi* Strains Isolated from Infected Foals in Poland. *POLOS ONE*, DOI: 10.1371/journal.pone.0152887.
- 22 Ilgekbayeva, GD, Otarbaev, BK, Kasymov, YI, Zanilabdin, MZ, Musaeva, GK, Kurman, S., Ilgekbayeva, AK. (2023). Biological characteristics of *Rhodococcus equi* isolated in the Republic of Kazakhstan. *Herald of Science of S.Seifullin Kazakh Agrotechnical Research University: Veterinary Sciences*, 2(002), 27-35. DOI: 10.51452/kazatuvc.2023.2.(002).1424.
- 23 Muscatello, G., Gilkerson, JR, Browning, GF. (2007). Comparison of two selective media for the recovery, isolation, enumeration and differentiation of *Rhodococcus equi*. *Veterinary Microbiology*, 119: 2–4, 324-329. DOI:10.1016/j.vetmic.2006.08.022.
- 24 Aphalo, PJ. (2020). *Learn R: as a language*. Chapman and Hall/CRC.
- 25 Ocampo-Sosa, AA, Lewis, DA, Navas, J. et al. (2007). Molecular epidemiology of *Rhodococcus equi* based on traA, vapA, and vapB virulence plasmid markers. *J Infect Dis.*, 196(5), 763–769;
- 26 Valero-Rello, A., Hapeshi, A., Anastasi, E. et al. (2015). An invertron-like linear plasmid mediates intracellular survival and virulence in bovine isolates of *Rhodococcus equi*. *Infect Immun.*, 83(7), 2725–2737.
- 27 Bryan, LK, Clark, SD, Díaz-Delgado, J. (2016). *Rhodococcus equi* infections in dogs. *Vet Path.*, 54(1), 159–163.
- 28 Flynn, O., Quigley, F., Costello, E., O’Grady, D., Gogarty, A., Mc Guirk, J., Takai, S. (2001). Virulence-associated protein characterisation of *Rhodococcus equi* isolated from bovine lymph nodes. *Vet. Microbiol.*, 78, 221–228.
- 29 von Bargen, K., Scraba, M., Krämer, I., Ketterer, M., Nehls, C., Krokowski, S. et al. (2019). Virulence-associated protein A from *Rhodococcus equi* is an intercompartmental pH-neutralising virulence factor. *Cellular microbiology*, 21(1), e12958.

References

- 1 Takai, S., Mizuno, Y., Suzuki, Y., Sasaki, Y., Kakuda, T., Kirikae, T. (2024). *Rhodococcus equi* infections in humans: an emerging zoonotic pathogen. *Nihon Saikingaku Zasshi*, 79(1), 15-24. DOI: 10.3412/jsb.79.15.
- 2 Val-Calvo, J., Darcy, J., Gibbons, J., Creighton, A., Egan, C., Buckley, T. et al. (2022). International Spread of Multidrug-Resistant *Rhodococcus equi*. *Emerging infectious diseases*, 28(9), 1899-903.
- 3 Rakowska, A., Cywinska, A., Witkowski, L. (2020). Current Trends in Understanding and Managing *Equine Rhodococcosis*. *Animals*, 10(10), 1910. DOI: 10.3390/ani10101910.
- 4 Lankenfeld, A., Weber, C., Rohn, K., Venner, M. (2021). Kinetics of serum amyloid A during the treatment period of foals with pneumonia. *Pferdeheilkunde – Equine Medicine*, 37.
- 5 Arnold-Lehna, D., Venner, M., Berghaus, LJ, Berghaus, R., Giguère, S. (2020). Changing policy to treat foals with *Rhodococcus equi* pneumonia in the later course of disease decreases antimicrobial usage without increasing mortality rate. *Equine veterinary journal*, 52(4), 531-7.
- 6 Bordin, AI, Huber, L., Sanz, MG, Cohen, ND. (2022). *Rhodococcus equi* foal pneumonia: Update on epidemiology, immunity, treatment and prevention. *Equine Vet J*, 54(3), 481-94.

- 7 Cohen, ND. (2014). *Rhodococcus equi* foal pneumonia. *Vet Clin North Am Equine Pract*, 30, 609–622.
- 8 Venner, M., Meyer-Hamme, B., Verspohl, J., Hatori, F., Shimizu, N., Sasaki, Y. et al. (2007). Genotypic characterization of VapA positive *Rhodococcus equi* in foals with pulmonary affection and their soil environment on a warmblood horse breeding farm in Germany. *Res Vet Sci.*, 83, 311–317.
- 9 Son, WG, Lee, DS, Yamatoda, N., Hatori, F., Shimizu, N., Kakuda, T. et al. (2006). Molecular typing of VapA-positive *Rhodococcus equi* isolates from Jeju native horses, Korea. *J Vet Med Sci.*, 68, 249–253.
- 10 Duquesne, F., Hébert, L., Sévin, C., Breuil, MF, Tapprest, J., Laugier, C. et al. (2010). Analysis of plasmid diversity in 96 *Rhodococcus equi* strains isolated in Normandy (France) and sequencing of the 87-kb type I virulence plasmid. *FEMS Microbiol Lett.*, 311, 76–81.
- 11 Takai, S., Chaffin, MK, Cohen, ND, Hara, M., Nakamura, M., Kakuda, T. et al. (2001). Prevalence of virulent *Rhodococcus equi* in soil from 5 R. equi-endemic horse-breeding farms and restriction fragment length polymorphisms of virulence plasmids in isolates from soil and infected foals in Texas. *J Vet Diagn Invest.*, 13, 489–494.
- 12 Sangal, V., Goodfellow, M., Jones, AL, Sutcliffe, IC. (2022). A stable home for an equine pathogen: valid publication of the binomial *Prescottella equi* gen. nov., comb. nov., and reclassification of four rhodococcal species into the genus *Prescottella*. *Int J Syst Evol Microbiol.*, 72(9).
- 13 Takai, S., Sekizaki, T., Ozawa, T., Sugawara, T., Watanabe, Y., Tsubaki, S. (1991). Association between a large plasmid and 15- to 17-kilodalton antigens in virulent *Rhodococcus equi*. *Infect Immun.*, 59, 4056–4060.
- 14 Takai, S., Shoda, M., Sasaki, Y., Tsubaki, S., Fortier, G., Pronost, S. et al. (1999). Restriction fragment length polymorphism of virulence plasmids in *Rhodococcus equi*. *J Clin Microbiol.*, 37, 3417–3420.
- 15 Takai, S., Son, WG, Lee, DS, Madarame, H., Seki, I., Yamatoda, N. et al. (2003). *Rhodococcus equi* virulence plasmids recovered from horses and their environment in Jeju, Korea: 90-kb type II and a new variant, 90-kb type. *V. J Vet Med Sci.*, 65, 1313–1317.
- 16 Takai, S., Suzuki, Y., Sasaki, Y., Kakuda, T., Ribeiro, MG, Makrai, L., Witkowski, L., Cohen, N., Sekizaki, T. (2023). Short review: Geographical distribution of equine-associated pVAPA plasmids in *Rhodococcus equi* in the world. *Vet Microbiol.*, 287, 109919. DOI: 10.1016/j.vetmic.2023.109919.
- 17 Takai, S., Anzai, T., Fujita, Y., Akita, O., Shoda, M., Tsubaki, S., Wada, R. (2000). Pathogenicity of *Rhodococcus equi* expressing a virulence-associated 20 kDa protein (VapB) in foals. *Veterinary Microbiology*, 76: 1, 71-80.
- 18 Ford, J., Santos-Aberturas, J., Hems, E., Sallmen, J., Bögeholz, L., Polturak, G., Osbourn, A., Wright, J., Rodnina, M., Vereecke, D., Francis, I., Truman, A. (2024). Discovery of lydiamycin A biosynthetic gene cluster in the plant pathogen *Rhodococcus fascians* guides structural revision and identification of molecular target. *BioRxiv*, DOI: 10.1101/2024.11.13.623425.
- 19 Suzuki, Y., Takahashi, K., Ishitsuka, T., Sugiyama, M., Sasaki, Y., Kakuda, T., Takai, S., Naito, I., Kohara, J. (2024). Experimental infection of goats with pVAPN-harboring *Rhodococcus equi* causes latent infection in the lymph nodes. *Vet Microbiol.*, 295, 110132. DOI: 10.1016/j.vetmic.2024.110132.
- 20 Suzuki, Y., Takahashi, K., Takase, F., Sawada, N., Nakao, S., Toda, A., Sasaki, Y., Kakuda, T., Takai, S. (2020). Serological epidemiological surveillance for vapN-harboring *Rhodococcus equi* infection in goats. *Comp Immunol Microbiol Infect Dis.*, 73, 101540. DOI: 10.1016/j.cimid.2020.101540.
- 21 Kalinowski, M., Grądzki, Z., Jarosz, Ł., Kato, K., Hieda, Y., Kakuda, T., Takai, S. (2016). Plasmid Profiles of Virulent *Rhodococcus equi* Strains Isolated from Infected Foals in Poland. *POLOS ONE*, DOI: 10.1371/journal.pone.0152887.
- 22 Ilgekbayeva, GD, Otarbaev, BK, Kasymov, YI, ZaniLabdin, MZ, Musaeva, GK, Kurman, S., Ilgekbayeva, AK. (2023). Biological characteristics of *Rhodococcus equi* isolated in the Republic of Kazakhstan. *Herald of Science of S.Seifullin Kazakh Agrotechnical Research University: Veterinary Sciences*, 2(002), 27-35. DOI: 10.51452/kazatuvc.2023.2.(002).1424.
- 23 Muscatello, G., Gilkerson, JR, Browning, GF. (2007). Comparison of two selective media for the recovery, isolation, enumeration and differentiation of *Rhodococcus equi*. *Veterinary Microbiology*, 119: 2–4, 324-329. DOI:10.1016/j.vetmic.2006.08.022.

- 24 Aphalo, P.J. (2020). *Learn R: as a language*. Chapman and Hall/CRC.
- 25 Ocampo-Sosa, AA, Lewis, DA, Navas, J. et al. (2007). Molecular epidemiology of *Rhodococcus equi* based on traA, vapA, and vapB virulence plasmid markers. *J Infect Dis.*, 196(5), 763–769;
- 26 Valero-Rello, A., Hapeshi, A., Anastasi, E. et al. (2015). An invertron-like linear plasmid mediates intracellular survival and virulence in bovine isolates of *Rhodococcus equi*. *Infect Immun.*, 83(7), 2725–2737.
- 27 Bryan, LK, Clark, SD, Díaz-Delgado, J. (2016). *Rhodococcus equi* infections in dogs. *Vet Path.*, 54(1), 159–163.
- 28 Flynn, O., Quigley, F., Costello, E., O’Grady, D., Gogarty, A., Mc Guirk, J., Takai, S. (2001). Virulence-associated protein characterisation of *Rhodococcus equi* isolated from bovine lymph nodes. *Vet. Microbiol.*, 78, 221–228.
- 29 von Barga, K., Scraba, M., Krämer, I., Ketterer, M., Nehls, C., Krokowski, S. et al. (2019). Virulence-associated protein A from *Rhodococcus equi* is an intercompartmental pH-neutralising virulence factor. *Cellular microbiology*, 21(1), e12958.

Herald of Science of S.Seifullin Kazakh Agro Technical Research University: Veterinary Sciences (Herald of Science KATRU: Veterinary Sciences)

GUIDELINES FOR AUTHORS

GENERAL

Herald of Science KATRU: Veterinary Sciences publishes peer-reviewed original articles and topical reviews on all aspects of veterinary sciences. Interdisciplinary articles and well-reasoned reviews are especially welcome.

The publication language is English. Publication of articles is free of charge. Submission of the manuscript is carried out through the Open Journal System online platform. Author registration at: <https://bulletinofscience.kazatu.edu.kz/index.php/veterinary-science>. Video instructions for registering authors are available at: <https://www.youtube.com/watch?v=UeZlKY4bozg>.

Herald of Science, KATRU: Veterinary Sciences uses double-blind peer review, which means the identities of the reviewer and author are hidden throughout the review process.

Before being sent to reviewers, manuscripts undergo a preliminary check to ensure compliance with the Journal's requirements. Criteria such as the novelty of the research, the originality of the results, compliance with the structure and format of the manuscript, the significance of the conclusions and the quality of the English language are checked.

Manuscripts that do not meet the requirements are rejected at this stage without review. Manuscripts are accepted with at least 70% text originality.

RESPONSIBILITY OF AUTHORS

By submitting a manuscript to *Herald of Science, KATRU: Veterinary Sciences*, the author(s) warrant that the manuscript is an original work and that has not been previously published or is not currently being considered for publication elsewhere. The authors also ensure that the ideas and words of others are properly acknowledged through appropriate attribution. Falsification of results and deliberately inaccurate statements, which may cause the manuscript to be rejected or the published article to be retracted, are not permitted.

The names of the authors should be listed in the manuscript in the order of their contribution to the article. Only those individuals who made a significant contribution to the study and preparation of the manuscript should be listed as authors. Those who assisted in the completion of the work, as well as the organizations that funded the research, are thanked in the "Acknowledgments" section at the end of the article.

The corresponding author must ensure that all co-authors have read and approved the final version of the article and agree to its publication. All authors of the manuscript are responsible for the information transmitted. The journal does not allow changes in authorship after prior acceptance.

REVIEWERS' RESPONSIBILITY

Reviewers must also register as a reviewer on the journal's website at: <https://bulletinofscience.kazatu.edu.kz/index.php/veterinary-science/user/register>.

The review must be submitted within two weeks. The ideas and hypotheses described in the manuscript should be kept confidential and not used for personal purposes. The results of the work cannot be discussed with other colleagues or used when conducting scientific seminars, giving lectures or presenting papers at scientific conferences.

Reviewers must clearly express their opinions, explaining and justifying their comments and be as objective as possible in assessing the manuscript. The reviewer's suggestions and advice for improving the work are welcome, even if he/she has decided to refuse publication.

Reviewers should pay attention to the novelty and/or originality of the work, the lack of similarity between the manuscript and other published articles, the completeness and correctness of the author(s) citing articles related to the topic (issue) being reviewed.

ETHICAL APPROVALS

Experiments using animals should be conducted in accordance with the principles established by the International Animal Ethics Committee or Institutional Ethics Committee and in accordance with local laws and regulations.

Research using animals must have approval from the Local Ethics Committee, which should be noted in the Materials and Methods section.

AUTHORS DECLARATION

The authors of the manuscript guarantee that the work and its main content have not been previously published and are not being considered for publication elsewhere. It should indicate the source of funding for the research (**download Sample of Author declaration form** <https://bulletinofscience.kazatu.edu.kz/index.php/veterinary-science/libraryFiles/downloadPublic/47>).

MANUSCRIPT TYPES ACCEPTED

- Original articles
- Review articles.

Editorial Board do not accept Case reports. The volume of original and/or review articles, excluding the Abstract and References, should not be less than 11,000 and 20,000 characters without spaces, respectively. The manuscript should not include texts, illustrations and tables that are not related to the study.

MANUSCRIPT FORMAT AND STRUCTURE

- Refer the latest articles and manuscript template from <https://bulletinofscience.kazatu.edu.kz/index.php/veterinary-science/demands> for format of each section (**download Manuscript template** <https://bulletinofscience.kazatu.edu.kz/index.php/veterinary-science/libraryFiles/downloadPublic/49>).

- Text should be printed observing the following margin sizes: top and bottom – 2 cm, left and right – 2 cm. Text alignment is widthwise (with automatic hyphenation). UDC is indicated in the upper left corner of the page.

- The manuscript must be typed in Times New Roman 12 point font, double spaced, in MS Word 2010.

- Lines must be numbered throughout the text.
- The title should be descriptive and not too long.

The title of the manuscript should be written in lowercase letters, except for the first letter of each main word. The manuscript should not include the name of the author(s) as it will undergo double-blind peer review. The title page indicating the full name and ORCID of the authors, affiliations, and e-mail are provided separately (**download Sample title page** <https://bulletinofscience.kazatu.edu.kz/index.php/veterinary-science/libraryFiles/downloadPublic/48>).

- **Abstract** should have the following structure: “Background and Aim”, “Materials and Methods”, “Results” and “Conclusion”, and should not exceed 300 words.

- **Key words** are given by semicolons in alphabetical order and in the singular (4-6 words and phrases).

- **Introduction:** a brief description of the relevance of the topic or issue based on a review of modern literature, justification of the significance of the aim, indicating the novelty of the work.

- **Materials and methods:** methods presented must be reproducible; the methods are briefly described without going into methodological features; For standard methods, reference to sources is required; When using a new method, a detailed description is required.
- **Results:** provides an analysis of the results obtained using tables, graphs and/or figures; the obtained data must be processed statistically.
- **Discussion:** a brief description of the most significant results of the work and comparison of the data obtained with the results of other studies; identifying regularities and/or contradictions.
- **Conclusion:** description of the conclusion and identification of further research on the topic (issue) under study.
- **Authors' Contributions, Acknowledgments and Conflicts of Interest** are included in a manuscript once it has been accepted for publication.
 - Author's Contributions: A brief summary of each author's contribution is provided.
 - Acknowledgment: information is provided about the funding of the work or words of gratitude are expressed to colleagues or other persons with whose assistance the research was carried out, etc.
 - Conflict of interest: Authors must confirm that there is no conflict of financial/non-financial interest.
 - **References:** In original and/or review articles, references from the last 5 years must be at least 40% and/or 50%, respectively. References should not include research reports, dissertations for academic degrees and conference proceedings.
 - References in the text should be given by placing in square brackets [1], [1,2,3]. Reference no. should start from no. 1 onwards from Introduction section. Reference list must be prepared simultaneously according to GOST 7.1-2003 (<https://bulletinofscience.kazatu.edu.kz/index.php/veterinary-science/gost>) and according to APA (<https://www.bibme.org/citation-guide/APA/book/>) with indicating the Digital Object Identifier (DOI), if available, in both cases.
 - Bibliography in other languages is described in the original according to GOST 7.1-2003 and transliterated into Latin in APA style using the link <http://translit-online.ru/>.

For example,

Description of the bibliography according to GOST 7.1-2003:

1 Думова, В.В. Противовирусные антитела в молозиве и молоке коров [Текст] / В.В. Думова, А.В. Мищенко, Т.Б. Никешина // Российский Ветеринарный журнал, - 2008.- Т.49.-С. 40-42.

Transliteration of bibliography using APA style:

1. Dumova, V.V., Mishchenko, A.V., Nikeshina, T.B. (2008). Protivovirusnye antitela v molozive i moloke korov [Antiviral antibodies in colostrum and milk of cows]. *Rossiiskii Veterinarnyi zhurnal [Russian Veterinary Journal]*, [In Russ.], 49: 40-42.

Authors for whom English is a second language should contact professional editors (e.g. ENAGO, Editage, etc.) and submit the manuscript along with an editing certificate. Alternatively, the manuscript may be edited by a native English speaker with a veterinary or medical background.

ABBREVIATIONS, TERMS AND NOMENCLATURE

Abbreviations of veterinary, medical and/or biological terms should only be used if they are generally known in world science. The full name of other terms must be included in parentheses when the abbreviation is first used.

The terms should be used in accordance with the latest recommendations of the relevant international nomenclature.

Bacterial names should conform to the latest edition of Burgee's Manual of Determinative Bacteriology (The Williams and Wilkins Co., Baltimore). Viruses must be classified and named as recommended by the International Committee on the Nomenclature of Viruses.

Names of microorganisms and zoological names should be in italics in the manuscript.

UNITS, SYMBOLS, TABLES, ILLUSTRATIONS AND FORMULAS

The units must be specified in accordance with the SI system. If symbols such as \times , μ , η , or v are used, they should be added using Word's symbol menu in Times New Roman.

The symbols such as ($^{\circ}$) or (\times) should be used from the symbol menu and not indicated by superscript letters: “o” or “x”. Spaces should be inserted between numbers and units of measurement (e.g., 3 kg), and between numbers and mathematical symbols (+, -, \times , =, <, >), but not between numbers and percentage symbols (e.g., 45%).

The table should be placed immediately after the text at the first mention. All tables in the manuscript must have links indicating their numbers (e.g., Table 1; Table 2, etc.). The title of the table should reflect its content, be accurate and concise. The name should be placed above the table on the left, without indentation (e.g., Table 1 - Hematological parameters of animals infected with *E. granulosus* protoscoleces). The table name is given without a dot at the end. If the table name takes up two or more lines, it should be written with one line spacing. Leave 1 space between the text and the figure before and after.

The illustrations (graphs, diagrams, diagrams, photographs and others) should be placed immediately after the text at the first mention. All illustrations in the manuscript must be referenced. When referring to a figure, the word “Figure” and its number should be written (e.g., In accordance with Figure 2...). The title of the figures should be made directly below the figure, aligned to the center. Leave 1 space between the text and the figure before and after.

The formulas. Simple in-line and one-line formulas must be typed in symbols without using special editors (the use of special symbols from the Symbol, GreekMathSymbols, MathPS, Math A Mathematica BTT fonts is allowed). Complex and multi-line formulas must be entirely typed in the formula editor Microsoft Equation 2.0, 3.0. It is not allowed to type part of the formula in symbols, and part in the formula editor.

REFERENCES

Each article must include bibliographic references. The name of the journal in which the cited article is published must be indicated as an abbreviated title from the cover of the corresponding journal, as well as via the link: *www.journalseek.net*, or from another verified list. The title of the journal should be written in italics.

MANUSCRIPT TRACKING

The corresponding author can track the progress of the submitted manuscript from his/her account. In addition, he/she will receive an email generated by the system (a certificate of the results of checking the text document for the presence of borrowings; a letter with comments from the editorial board and/or reviewers; a letter from the editorial board of the journal on the acceptance and/or rejection of the manuscript, etc.).

MANUSCRIPT REVIEW, PROOF READING AND PUBLICATION

Manuscript review. The corresponding author will receive an email once reviewers submit comments to *Herald of Science, KATRU: Veterinary Sciences*. The Editor will check the reviewers' comments and forward them to the corresponding author with or without additional comments. The corresponding author must submit an edited manuscript based on comments

from the editors and/or reviewers within 4 weeks. If the corresponding author requires more time, he/she must obtain the editor's approval. If the author does not agree with the reviewer's comments, he/she provides a rationale for each comment. The final decision on each manuscript is made by the Editor-in-chief.

Proof corrections. The corresponding author will receive an e-mail containing PDF. The proofread text must be returned to the Editor within three days of receipt with corrections, if any. If there are no corrections, the corresponding author should respond accordingly. The author(s) are responsible for all comments made on their work, including any changes made.

All comments/questions from the Editor and reviewers, as well as corrections and responses from the corresponding author, must be reflected in the same text of the manuscript using Track changes.

Publication. Articles published online and/or in paper version of the *Herald of Science KATRU: Veterinary Sciences* are assigned a DOI number.

MANUSCRIPT STRUCTURE

UDC

Title of the manuscript submitted to the Journal

(The title of the manuscript should be written in lowercase letters, except for the first letter)

Abstract

Background and Aim

Materials and Methods.....

Results.....

Conclusion.....(max. 300 words)

Key words:(4-6 words)

Introduction

.....

Materials and methods

.....

Results

.....

Discussion

.....

Conclusion

.....

Authors' Contributions

.....

Acknowledgements

.....

Conflicts of Interest

.....

References

.....

SAMPLE TITLE PAGE

UDC 619.616.98.579.841.935.07

Using indirect hemagglutination assay for the diagnosis of cattle brucellosis

Mikhail M. Mikailov¹ , Shahrudin A. Gunashev¹ ,
Elmira A. Yanikova¹ , Ahmed A. Halikov¹ , Aitbay. K. Bulashev² 

¹Laboratory of Infectious Pathology of Farm Animals, Caspian Zonal Research
Veterinary Institute, Republic of Dagestan, Makhachkala, Russia;

²Department of Microbiology and Biotechnology,
Faculty of Veterinary Medicine and Livestock Technology,
Saken Seifullin Kazakh Agrotechnical Research University, Astana, Kazakhstan.

Corresponding author: Aitbay K. Bulashev: aytbay57@mail.ru

Co-authors: (1:MM)* mikail.mikailov1981@mail.ru; (2:ShG) sgunashev@mail.ru;
(3:EY) nvetmedservis@mail.ru; (4:AH) axmedx93@mail.ru

ORCID:

1st Author: <https://orcid.org/0000-0002-9620-431X>

2nd Author: <https://orcid.org/0000-0003-4804-2755>

3rd Author: <https://orcid.org/0000-0002-5561-2499>

4th Author: <https://orcid.org/0000-0002-9765-008X>

5th Author: <https://orcid.org/0000-0002-8427-509X>

*Note: Abbreviations of author names should be used for “Authors’ Contributions”, e.g.:
“Authors’ Contributions

AB and MM: Conceptualized and designed the study, conducted a comprehensive literature search, analyzed the gathered data and drafted the manuscript. ShG, EY and AH: Conducted the final revision and proofreading of the manuscript. All authors have read, reviewed, and approved the final manuscript”.

AUTHOR DECLARATION
Herald of Science, KATRU: Veterinary Sciences

We, the undersigned, author(s) of the manuscript entitled _____

_____ hereby declare

that:

- the manuscript is our own original work, it has not been published before and is not currently being considered for publication elsewhere;
- the research results, ideas and/or words of others have been properly cited through appropriate references and/or citations, and accurately referenced;
- we take full responsibility for the information presented in the manuscript as our own results;
- the research results and conclusions are reliable and presented sufficiently fully, correctly and objectively;
- the authors of the manuscript include only persons who have made a significant contribution to obtaining research results;
- permission of ethical commission has been taken and included a statement in the manuscript;
- we have no competing interests;
- the final version of the manuscript has been reviewed, read and approved for publication in Herald of Science, KATRU: Veterinary Sciences.

Full name and address of funding source

We also agree to attribution in the following order:

Author's name	Signature	e-mail, mobile phone number
1. _____	_____	_____
2. _____	_____	_____
3. _____	_____	_____
4. _____	_____	_____
5. _____	_____	_____
6. _____	_____	_____
7. _____	_____	_____
8. _____	_____	_____

Date: _____

Corresponding author address, e-mail and mobile phone number: _____

**Name, signature and seal of
 Head of Department/Institute**

CONTENT

VETERINARY SCIENCES

<i>O. S. Akibekov, F. Tosini, A. S. Syzdykova, A. M. Gajimuradova, D. D. Zhumalina, F. S. Zhagipar, N.N. Gubaidullin, N. A. Askarova, S. S. Yesseneyeva</i>	4
Study of the Immunologic Value of <i>Trichinella spiralis</i> recombinant Protein Tsp-LE.....	
<i>Sh. K. Sabyrbekova, G. D. Ilgekbayeva</i>	16
Meat Quality Control in Beef Production.....	
<i>R. M. Rychshanova, A.M. Mendybayeva, Y. Y. Aleshina, M. S. Ernazarova, Z. B. Abilova, A. R. Gabitova</i>	34
Analysis of microbial contamination in different types of meat in the Kostanay region.....	
<i>S. T. Dyusembayev, A. T. Serikova, Sh. K. Suleimenov, Zh. T. Serikov, Zh. Sh. Adilbekov</i>	43
Plants with radioprotective properties: current research and application prospects.....	
<i>G. D. Ilgekbayeva, Y.I. Kassymov, B. K. Otarbayev, M.Z. Zaniabdin, S. Kurman, N. Matikhan, U. Zholdas, B. A. Valieva</i>	51
Antigenic Activity of Various <i>Rhodococcus equi</i> Strain Plasmids.....	

HERALD OF SCIENCE

*S. Seifullin Kazakh Agrotechnical Research
University*

№ 4 (008) 2024

Journal is registered with the Information and Archives Committee,
Ministry of Culture, Information and Sports, Republic of Kazakhstan
(Certificate №5770-Ж)
(Certificate №13279-Ж)

Editor N.K. Kokumbekova
Technical Editor M.M. Zhumabekova
Computer layout S.S. Romanenko

Submitted for printing 10.09.2024, Signed for printing 26.12.2024,
Format 60×84 ¹/₈, Conditionally printed sheets 8,5, Journal circulation 300,
Order № 24156

Printed in the Publishing House of S. Seifullin Kazakh
Agrotechnical Research University
010011, Zhenis avenue 62a, Astana, Kazakhstan
Phone: +7-7172-31-02-75